

LITERATUR

MOLLUSCA (WEICHTIERE)

ARTIKEL ZU GASTROPODA (SCHNECKEN)

- BLANCHARD, M. (1997): Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe. Current state and consequences. *Sci. Mar.* 61, Suppl 2: 109-118
- FILIPPENKO, D.P. (2011): Invasive Species *Potamopyrgus antipodarum* (Gray, 1843) (Mollusca: Gastropoda: Hydrobiidae) in the waterbodies of Kaliningrad Region (Russia). *Russ J Biol Invasions* 2: 286-292
- FILIPPENKO, D.P. & SON, M.O. (2008): The New Zealand mud snail *Potamopyrgus antipodarum* (Gray, 1843) is colonizing the artificial lakes of Kaliningrad City, Russia (Baltic Sea Coast). *Aquatic Invasions* 3: 345-347
- MINCHIN, D., MCGRATH, D. & DUGGAN, C.B. (1995): The Slipper Limpet, *Crepidula fornicata* (L.), in Irish waters, with a review of its occurrence in the North-Eastern Atlantic. *J. Conchol.*, London 35: 247-256
- PONDER, W.F. (1988): *Potamopyrgus antipodarum* – a molluscan coloniser of Europe and Australia. *J Moll Stud* 54: 271-285
- THIELTGES, D.W. (2005): Benefit from an invader: American slipper limpet *Crepidula fornicata* reduces star fish predation on basibiont European mussels. *Hydrobiologia* 541: 241-244
- THIELTGES, D.W., STRASSER, M. & REISE, K. (2003): The American slipper limpet *Crepidula fornicata* (L.) in the northern Wadden Sea 70 years after its introduction. *Helgol Mar Res* 57: 27-33
- THIELTGES, D.W., STRASSER, M., VAN BEUSEKOM, J. & REISE, K. (2004): Too cold to prosper – winter mortality prevents population increase of the introduced slipper limpet *Crepidula fornicata* in northern Europe. *J Exp Mar Biol Ecol* 311: 375-391
- THIELTGES, D.W., REISE, K., PRINZ, K. & JENSEN, K.T. (2009): Invaders interfere with native parasite-host interactions. *Biol Invasions* 11: 1421-1429

ARTIKEL ZU BIVALVIA (MUSCHELN)

- ARMONIES, W. & REISE, K. (1999): On the population development of the introduced razor clam *Ensis americanus* near the island of Sylt (North Sea). *Helgoländer Meeresunters* 52: 291-300
- BÄTKE, J. (1994): Die Verbreitung von *Corbicula fluminalis* (O.F. Müller 1774) (Bivalvia, Corbiculidae) in der Weser. *Lauterbornia* 15: 17-21
- BAMBER, R.N. & TAYLOR, J.D. (2002): The brackish water mussel *Mytilopsis leucophaeta* (Conrad, 1831) (Bivalvia: Dreissenidae) in the river Thames. *J Conch* 37: 403-404

- BEHREND, B., HERTWECK, G., LIEBEZEIT, G. & GOODFRIEND, G. (2005): Earliest Holocene occurrence of the soft-shell clam, *Mya arenaria*, in the Greifswalder Bodden, Southern Baltic. *Marine Geology* 216: 79-82
- BEUKEMA, J.J. & DEKKER, R. (1995): Dynamics and growth of a recent invader into European coastal waters: the American razor clam, *Ensis directus*. *J mar biol Ass U.K.* 75: 351-362
- BIRNBAUM, C. (2011): NOBANIS – Invasive Alien Species Fact Sheet – *Dreissena Polymorpha*. From: Online Database of the European Network on Invasive Alien Species – NOBANIS www.nobanis.org (20.03.2013)
- BOCK, G., LIEBERUM, C., SCHÜTT, R., WIESE, V. (2015): Erstfund der Brackwassermuschel *Rangia cuneata* in Deutschland. *Schr Malakozool* 28: 13-16
- BOETTGER, C.R. (1933a): Über die Ausbreitung der Muschel *Congeria cochleata* Nyst in europäischen Gewässern und ihr Auftreten im Nordostseekanal. *Zool Anz* 101: 43-48
- BOUDRY, P., HEURTEBISE, S., COLLET, B., CORNETTE, F. & GÉRARD, A. (1998): Differentiation between populations of the Portuguese oyster, *Crassostrea angulata* (Lamarck) and the Pacific oyster, *Crassostrea gigas* (Thunberg), revealed by mtDNA RFLP analysis. *J Exp Mar Biol Ecol* 226: 279-291
- CROSS, M.E., BRADLEY, C.R., CROSS, T.F., CULLOTY, S., LYNCH, S., MCGINNITY, P., O'RIORDAN, R.M., VARTIA, S. & PRODÖHL, P.A. (2016): Genetic evidence supports recolonization by *Mya arenaria* of western Europe from North America. *Mar Ecol Prog Ser* 549: 99-112
- DARR, A. & ZETTLER, M.L. (2000): Erstnachweis von *Congeria leucophaeta* (Conrad, 1831) in Mecklenburg-Vorpommern (Bivalvia: Dreissenidae). *Malakologische Abhandlungen* 20: 197-200
- DIEDERICH, S., NEHLS, G., VAN BEUSEKOM, J.E.E. & REISE, K. (2005): Introduced Pacific Oysters (*Crassostrea gigas*) in the northern Wadden Sea: invasion accelerated by warm summers? *Helgol Mar Res* 59: 97-106
- DOLMER, P., HOLM, M.W., STRAND, Á., LINDEGARTH, S., BODVIN, T., NORLING, P. & MORTENSEN, S. (2014): The invasive Pacific oyster, *Crassostrea gigas*, in Scandinavian coastal waters: A risk assessment on the impact in different habitats and climate conditions. *Fiske og Havet, Havforskningsinstituttet*, nr. 2/2014, 67 pp
- DZIUBINSKA, A. (2011): *Mytilopsis leucophaeata*, an alien dreissenid bivalve discovered in the Gulf of Gdansk (southern Baltic Sea). *Oceanologia* 53: 651-655
- ESCHWEILER, N. & CHRISTENSEN, H.T. (2011): Trade-off between increased survival and reduced growth for blue mussels living on Pacific oyster Reefs. *J Exp Mar Biol Ecol* 403: 90-95
- FLORIN, A.-B., MO, K., SVENSSON, F., SCHAGERSTRÖM, E., KAUTSKY, L. & BERGSTRÖM, L. (2013): First records of Conrad' false mussel, *Mytilopsis leucophaeta* (Conrad, 1831) in the southern Bothnian Sea, Sweden, near a nuclear power plant. *BiolInvasions Records* 2: 303-309
- FORSSTRÖM, T., FOWLER, A.E., LINDQUIST, M. & VESAKOSKI O. (2016): The introduced dark false mussel, *Mytilopsis leucophaeata* (Conrad, 1831) has spread in the northern Baltic Sea. *BiolInvasions Records* 5: 81-84

- GROSLIER, T., CHRISTENSEN, H.T., DAVIDS, J., DOLMER, P., ELMEDAL, I., HOLM, M.W. & HANSEN, B.W. (2014): Status of the Pacific Oyster *Crassostrea gigas* (Thunberg, 1793) in the western Limfjord, Denmark – Five years of population development. *Aquatic Invasions* 9: 175-182
- GÜRS, I., GÜRS, K., PAUSTIAN, J. & WIESE, V. (1993): *Ensis directus* (Conrad, 1843) zum ersten Mal in der südwestlichen Ostsee (Bivalvia: Solenidae). *Schriften aus dem Malakozoologischen Haus der Natur* 6: 79
- HOLM, M.W., DAVIDS J.K., DOLMER P., VISMANN B., HANSEN B.W. (2015): Moderate establishment success of Pacific oyster, *Crassostrea gigas*, on a sheltered intertidal mussel bed. *J Sea Res* <http://dx.doi.org/10.1016/j.seares.2015.07.009>
- HOLM, M. W., DAVIDS, J.K., DOLMER, P., HOLMES, E., NIELSEN, T.T., VISMAN, B. & HANSEN, B. W. (2015): Coexistence of Pacific oyster *Crassostrea gigas* (Thunberg, 1793) and blue mussels *Mytilus edulis* Linnaeus, 1758 on a sheltered intertidal bivalve bed? *Aquatic Invasions* 11: 155-165
- HOPPE, K.N. (2002): *Teredo navalis* – the cryptogenic shipworm. In: LEPPÄKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic Species of Europe. Distribution, Impacts and Management*. Kluwer Academic Publishers, Dordrecht, Boston, London, 116-119
- HUVET, A., GÉRARD, A., LEDU, C., PHÉLIPOT, P., HEURTEBISE, S. & BOUDRY, P. (2002): Is fertility of hybrids enough to conclude that the two oysters *Crassostrea gigas* and *Crassostrea angulata* are the same species? *Aquat Living Resour* 15: 45-52
- IMO, M, SEITZ, A.. & JOHANNESSEN, J. (2010): Distribution and invasion genetics of the quagga mussel (*Dreissena rostriformis bugensis*) in German rivers. *Aquat Ecol* 44: 731-740
- JENSEN, K.R. (2010a): NOBANIS – Invasive Alien Species Fact Sheet – *Petricola Pholadiformis*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (15.03.2013)
- JUEG, U. & ZETTLER, M. (2004): Die Molluskenfauna der Elbe in Mecklenburg-Vorpommern mit Erstnachweis der Grobgerippten Körbchenmuschel *Corbicula fluminea* (O.F. Müller 1756). *Mitteilungen der NGM*, 4: 85-89
- KENNEDY, V.S. (2011): The invasive dark falsemussel *Mytilopsis leucophaeata* (Bivalvia: Dreissenidae): a literature review. *Aquat Ecol* 45: 163-183
- LAINE, A.O., MATTILA, J. & LEHIKONEN, A. (2006): First record of the brackish water dreissenid bivalve *Mytilopsis leucophaeata* in the northern Baltic Sea. *Aquatic Invasions* 1: 38-41
- LUCZAK, C., DEWARUMEZ, J.-M. & ESSINK, K. (1993): First Record of the American Jack Knife Clam *Ensis directus* on the French Coast off the North Sea. *J. mar. biol. Ass.* 73: 233-235
- MEIXNER, R. (1984): Erster Wildfund einer Pazifischen Auster *Crassostrea gigas*. *Inf Fischwirt* 31: 140
- MEßNER U. & ZETTLER M. (2015): Die Quagga-Muschel *Dreissena (Pontodreissena) bugensis* (Andrusov, 1897) hat die Mecklenburgische Seenplatte und das Oderhaff erreicht (Bivalvia: Dreissenidae). *Lauterbornia* 80: 31-35

- MEYER-WAARDEN, P.F. (1964): Mastversuche mit portugiesischen Austern im deutschen Wattengebiet. *Inf Fischwirt* 11: 77-78
- MINCHIN, D., LUCY, F. & SULLIVAN, M. (2002): Zebra Mussel: Impacts and Spread. In: LEPPÄKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, Boston, London, 135-146
- MIOSSEC, L., LE DEUFF, R.-M. & GOULLETQUER, P. (2009): Alien species alert: *Crassostrea gigas* (Pacific oyster). ICES Cooperative Research Report No. 299
- MOEHLER, J., WEGNER, K.M., REISE, K. & JACOBSEN, S. (2011): Invasion genetics of Pacific oyster *Crassostrea gigas* shaped by aquaculture stocking practices. *J Sea Res* 66: 256-262
- NEHRING, S. (2011): NOBANIS – Invasive Alien Species Fact Sheet – *Crassostrea gigas*. From: Online Database of the North European and Baltic Network on Invasive Alien Species. NOBANIS www.nobanis.org (01.03.2013)
- NEUDECKER, T. (1990): The history of the former German oyster fishery and mariculture: 400 Years of Crown Law on oysters (4.2.1587). *Dt Hydrograph Zeitschr, Ergänzungsheft B* 22
- OLIVER, P.G., HOLMES, A.M. & METTAM, C. (1998): *Mytilopsis leucophaeta* (Conrad, 1831) (Bivalvia: Dreissenoidea). A species new to the British fauna. *J Conch* 36: 13-18
- PERNET, F., BARRET, J., LE GALL, P., CORPOREAU, C., DÉGREMONT, L., LAGARDE, F., PÉPIN, J.-F. & KECK, N. (2012): Mass mortalities of Pacific Oysters *Crassostrea gigas* reflect infectious diseases and vary with farming practices in the Mediterranean Thau lagoon, France. *Aquacult Environm Interact* 2: 215-237
- PETERSEN, K.S., RASMUSSEN, K.L., HEINEMEIER, J. & RUD, N. (1992): Clams before Columbus? *Nature* 359: 679
- PFENNINGER, M., REINHARDT, F. & STREIT, B. (2002): Evidence for cryptic hybridization between different evolutionary lineages of the invasive clam genus *Corbicula* (Veneroidea, Bivalvia). *J Evol Biol* 15: 818-829
- RAYBAUD, V., BEAUGRAND, G., DEWARUMEZ, J.-M. & LUCZAK C. (2015): Climate induced range shifts of the American jackknife clam *Ensis directus* in Europe. *Biol Invasions* 17: 725-741
- REISE, K. (1998): Pacific Oysters invade mussel beds in the European Wadden Sea. *Senckenbergiana marit*, 28: 167-175
- RUDINSKAYA L.V. & GUSEV A.A. (2012): Invasion of the North American Wedge Clam *Rangia cuneata* (G.B. Sowerby I, 1831) (Bivalvia: Mactridae) in the Vistula Lagoon of the Baltic Sea. *Russ J Biol Invas* 3: 220-229
- SCHLESCH, H. (1932): Über die Einwanderung nordamerikanischer Meeresmollusken in Europa unter Berücksichtigung von *Petricola pholadiformis* Lam. und ihre Verbreitung im dänischen Gebiet. *Arch Moll* 64: 146-154
- SCHÜTZ, L. (1961): Verbreitung und Verbreitungsmöglichkeiten der Bohrmuschel *Teredo navalis* L. und ihr Vordringen in den NO-Kanal bei Kiel. *Kieler Meeresforsch.* 17: 228-236

- SCHWEIMANN, M. (1993): Biodeterioration durch holzerstörende Organismen entlang der Küsten Schleswig-Holsteins. *Schr Malakozool Cismar* 6: 6-8
- SMITH, P., GUY, C. & DONNAN D. (2015): Pacific Oysters, *Crassostrea gigas*, established in Scotland. *Aquatic Conserv Mar Freshw Ecosyt* 25: 733-742
- SORDYL, H., BÖNSCH, R., GERCKEN, J., GOSSELCK, F., KREUZBERG, M. & SCHULZE, H. (1998): Verbreitung und Reproduktion des Schiffsbohrwurms *Teredo navalis* L. an der Küste Mecklenburg-Vorpommerns. *Dt Gewässerkd Mitt* 42: 142-149
- STRAND, A., BLANDA, E., BODVIN, T., DAVIDS, J.K., JENSEN, L.F., HOLM-HANSEN, T.H., JELMERT, A., LINDEGARTH, S., MORTENSEN, S., MOY, F.E., NIELSEN, P., NORLING, P., NYBERG, C., CHRISTENSEN, H.T., VISMANN, B., HOLM, M.W., HANSEN, B.W. & DOLMER, P. (2012): Impact of an icy Winter on the Pacific oyster (*Crassostrea gigas* Thunberg, 1793) populations in Scandinavia. *Aquatic Invasions* 7: 433-440
- STRASSER, M. (1999): *Mya arenaria* – an ancient invader of the North Sea coast. *Helgoländer Meeresunters* 52: 309-324
- TITTIZER, T. & TAXACHER, M. (1997): Erstnachweis von *Corbicula fluminea/fluminalis* (Müller 1774) (Corbiculidae, Mollusca) in der Donau. *Lauterbornia* 31: 103-107
- TROOST, K. (2010): Causes and effects of a highly successful marine invasion: Case-study of the introduced Pacific oyster *Crassostrea gigas* in continental NW European estuaries. *J Sea Res* 64: 145-165
- TUENTE, U., PIEPENBURG, D. & SPINDLER, M. (2002): Occurrence and settlement of the common shipworm *Teredo navalis* (Bivalvia: Teredinidae) in Bremerhaven harbours, northern Germany. *Helgol Mar Res* 56: 87-94
- VERWEEN, A., VINCX, M., MEES, J. & DEGRAER, S. (2005): Seasonal variability of *Mytilopsis leucophaeata* larvae in the harbour of Antwerp: implications for ecologically and economically sound biofouling control. *Belg J Zool* 135: 91-93
- VERWEEN A., KERCKHOF F., VINCX M., DEGRAER S. (2006): First European record of the invasive brackish water clam *Rangia cuneata* (G.B. Sowerby I, 1831) (Mollusca: Bivalvia). *Aquatic Invasions* 1: 198-203
- VOIGT, M. (1999): *Ensis directus*. In: GOLLASCH, S., MINCHIN, D., ROSENTHAL, H. & VOIGTS, M. (Hrsg.): Exotics across the ocean. Case histories on introduced species: their general biology, distribution, range expansion and impact. Kiel 1999, 37-42
- VON COSEL, R., DÖRJES, J. & MÜHLENHARDT-SIEGEL, U. (1982): Die amerikanische Schwertmuschel *Ensis directus* (Conrad) in der Deutschen Bucht. 1. Zoogeographie und Taxonomie im Vergleich mit den einheimischen Schwertmuschel-Arten. *Senckenbergiana marit* 14: 147-173
- WANG, H., QIAN, L., LIU, X., ZHANG, G. & GUO, X. (2010): Classification of a common cupped oyster from southern China. *J Shelf Res* 29: 857-866
- WATERMANN, B.T., HERLYN, M., DAEHNE, B., BERGMANN, S., MEEMKEN, M. & KOLODZEY, H. (2008): Pathology and mass mortality of Pacific oysters, *Crassostrea gigas* (Thunberg), in 2005 at the East Frisian coast, Germany. *J Fish Dis* 31: 621-630

WEHRMANN, A., HERLYN, M., BUNGENSTOCK, F., HERTWECK, G. & MILLAT, G. (2000): The distribution gap is closed – first record of naturally settled Pacific Oysters *Crassostrea gigas* in the East Frisian Wadden Sea, North Sea. *Senckenbergiana marit.* 30: 153-160

WOLFF, W.J. & REISE, K. (2002): Oyster imports as a vector for the introduction of alien species into northern and western European coastal waters. In: LEPPAKÖSKI, E., OLENIN, S. & GOLLASCH, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management.* Kluwer Academic Publishers, Dordrecht, Boston, London, 193-205

WOŹNICZKA A., WAWRZYNIAK-WYDROWSKA B., RADZIEJEWSKA T. & SKRZYPACZ A. (2016): The quagga mussel (*Dreissena rostriformis bugensis* Andrusov, 1897) - another Ponto-Caspian dreissenid bivalve in the southern Baltic catchment: the first record from the Szczecin Lagoon. *Oceanologia* (2015), <http://dx.doi.org/10.1016/j.oceano.2015.12.002>

WRANGE, A.-L., VALERO, J., HARKESTAD, L.S., STRAND, O., LINDEGARTH, S., CHRISTENSEN, H.T., DOLMER, P., KRISTENSEN, P.S. & MORTENSEN, S. (2010): Massive settlements of the Pacific oyster, *Crassostrea gigas*, in Scandinavia. *Biol Invasions* 12: 1453-1458

ÜBERSICHTSARTIKEL UND LISTEN MIT ANGABEN ZU GASTROPODA UND BIVALVIA

ENO, N.C., CLARK, R.A. & SANDERSON, W.G. (eds.) (1997): *Non-native marine species in British waters: a review and directory.* Joint Nature Conservation Committee, Peterborough, 152 pp

GLÖER, P. (2002): Die Süßwassergastropoden Nord- und Mitteleuropas, 327 pp. In: *Die Tierwelt Deutschlands* (F. Dahl), 73. Teil, ConchBooks Hackenheim

GOLLASCH, S. & NEHRING, S. (2006): National checklist for aquatic species in Germany *Aquatic invasions* 1: 245–269

GOLLASCH, S., HAYDAR, D., MINCHIN, D., WOLFF, W.J. & REISE, K. (2009): Introduced Aquatic Species of the North Sea Coasts and Adjacent Brackish Waters. In: RILOV, G. & CROOKS, J.A. (eds.): *Biological invasions in marine ecosystems*, Chapter 29: 507-528 *Ecological Studies* 204, Springer-Verlag Berlin Heidelberg 2008

GOULLETQUER, P., BACHELET, G., SAURIAU, P.G. & NOEL, P. (2002): Open atlantic coast of Europe – a century of introduced species into French waters. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management.* Kluwer Academic Publishers, Dordrecht, Boston, London, 276-290

HAYWARD, P.J. & RYLAND, J.S. (1990): *The Marine fauna of the British Isles and North-West Europe*, Vol. 1 & 2. Oxford Science Publications, Clarendon Press, Oxford, 1990

JENSEN, K.T. (1992): Macrozoobenthos on an intertidal mudflat in the Danish Wadden Sea: comparisons of surveys made in the 1930s, 1940s and 1980s. *Helgoländer Meeresunters.* 46: 363-376

JENSEN, K. & KNUDSEN, J. (2005): A summary of alien marine benthic invertebrates in Danish waters. *Oceanol Hydrobiol Stud* 34, Suppl 1: 137-162

- KERCKHOF, F., HAELTERS, J. & GOLLASCH, S. (2007): Alien species in the marine and brackish water ecosystems: the situation in Belgian waters. *Aquatic Invasions* 2: 243-257
- LACKSCHEWITZ, D., REISE, K. & BUSCHBAUM, C. (2012, 2013): Neobiota. In: Monitoring und Bewertung des Benthos, der Lebensraumtypen/Biotope und der Gebietsfremden Arten (Cluster 4, Benthosmonitoring, Synthetischer Jahresbericht)
- MINCHIN, D., COOK, E.J. & CLARK, P.F. (2013): Alien species in British brackish and marine waters. *Aquatic Invasions* 8, Suppl: 3-19
- NEHRING, S. (2000a): Neozoen im Makrozoobenthos der deutschen Ostseeküste. *Lauterbornia* 39: 117-126
- NEHRING, S. (2006): Four arguments why so many alien species settle into the estuaries, with special reference to the German river Elbe. *Helgol Mar Res* 60: 127-134
- NEHRING, S. & LEUCHS, H. (1999a): Neozoa (Makrobenthos) an der deutschen Nordseeküste. Eine Übersicht. Bundesanstalt für Gewässerkunde, Koblenz. 131 pp
- POST, D. & LANDMANN, M. (1994): Verbreitungsatlas der Fließgewässerfauna in Ostfriesland. Staatl Amt Wasser Abfall, 141 S.
- REISE, K. (1982): Long-term changes in the macrobenthic invertebrate fauna of the Wadden Sea: are polychaetes about to take over? *Neth J Sea Res* 16: 29-36
- REISE, K., HERRE, E. & STURM, M. (1994): Biomass and abundance of macrofauna in intertidal sediments of Königshafen in the northern Wadden Sea. *Helgoländer Meeresunters.* 48: 201-215
- REISE, K., DANKERS, N. & ESSINK, K. (2005): Introduced species. In: ESSINK, K, DETTMANN, C., FARKE, H., LAURSEN, K., LÜERßEN, G., MARENCIC, H. & WIERSINGA, W. (eds.): *Wadden Sea Quality Status Report 2004. Wadden Sea Ecosystem No. 19. Common Wadden Sea Secretariat, Wilhelmshaven, Germany, 155-161*
- TITTIZER, T., SCHÖLL, F., BANNING, M., HAYBACH, A. & SCHLEUTER, M. (2000): Aquatische Neozoen im Makrobenthos der Binnenwasserstraßen Deutschlands. *Lauterbornia* 39: 1-72
- WITT, J. & KRUMWIEDE, A. (2006): Entwicklung eines Bewertungskonzeptes und Bewertung des Makrozoobenthos der Unterweser. Anhang: Beschreibung und Bewertung des Ist-Zustandes. Im Auftrag des Wasser- und Schifffahrtsamtes Bremerhaven (WSA)
- WOLFF, W.J. (2005): Non-indigenous marine and estuarine species in The Netherlands. *Zool Meded* 79: 1-116

Listen und Internetquellen zu Mollusca

www.aquatic-aliens.de
www.frammandearter.se