

LITERATUR

ANNELIDA

ARTIKEL ZU POLYCHAETA (VIELBORSTER)

- AMBROSE, W.G. (1984): Influences of predatory polychaetes and epibenthic predators on the structure of a soft-bottom community in a Maine estuary. *J Exp Mar Biol Ecol* 81: 115-145
- ATKINS, S.M., JONES, A.M. & GARWOOD, P.R. (1987): The ecology and reproductive cycle of a population of *Marenzelleria viridis* (Annelida: Polychaeta: Spionidae) in the Tay Estuary. *Proc Royal Soc Edinburgh*, 92B: 311-322
- BASTROP, R. & BLANK, M. (2006): Multiple invasions – a polychaete genus enters the Baltic Sea. *Biol Invasions* 8: 1195-1200
- BAZTERRICA, M.C., BOTTO, F. & IRIBARNE, O. (2012): Effects of an invasive reef-building polychaete on the biomass and composition of estuarine macroalgal assemblages. *Biol Invasions* 14: 765-777
- BICK, A. & GOSSELCK, F. (1985): Arbeitsschlüssel zur Bestimmung der Polychaeten der Ostsee. *Mitt zool Mus Berl.* 61: 171-272
- BICK, A. & BURCKHARDT, R. (1989): Erstnachweis von *Marenzelleria viridis* (Polychaeta Spionidae) für den Ostseeraum, mit einem Bestimmungsschlüssel der Spioniden der Ostsee. *Mitt zool Mus Berlin* 65: 237-247
- BLAKE, J.A. (1991): Revision of some genera and species of Cirratulidae (Polychaeta) from the Western North Atlantic. *Ophelia*, Suppl 5: 17-30
- BLAKE, J.A., GÖRANSSON P. (2015): Redescription of *Thary killariensis* (Southern) from Ireland and description of two new species of *Tharyx* from the Kattegat, Sweden (Polychaeta, Cirratulidae). *Zootaxa* 4039: 501-515
- BLANK, M., BASTROP, R., RÖHNER, M. & JÜRSS, K. (2004): Effect of salinity on spatial distribution and cell volume regulation in two sibling species of *Marenzelleria* (Polychaeta: Spionidae). *Mar Ecol Prog Ser* 271: 193-205
- BLANK, M., LAINE, A.O., JÜRSS, K. & BASTROP, R. (2008): Molecular identification key based on PCR/RFLP for three polychaete sibling species of the genus *Marenzelleria*, and the species' current distribution in the Baltic Sea. *Helgol Mar Res* 62: 129-141
- DAUER, D.M., MAHON H.K., SARDÁ R. (2003): Functional morphology and feeding behavior of *Streblospio benedicti* and *S. shrubsolii*. *Hydrobiologia* 496: 207-213
- DEAN, D. (1978): Migration of the sandworm *Nereis virens* during winter nights. *Mar Biol* 45: 165-173
- DIDŽIULIS, V. (2006): NOBANIS – Invasive Alien Species Fact Sheet – *Marenzelleria neglecta*. From: Online Database of the North European and Baltic Network on Invasive Alien Species – NOBANIS www.nobanis.org (29.05.2012)

- DITTMER, J.D. (1981): The distribution of subtidal macrobenthos in the estuaries of the rivers Ems and Weser In: DANKERS, N., KÜHL, H. & WOLFF, W. (Hrsg.): Invertebrates of the Wadden Sea, Report 4. Balkema Rotterdam, 188-206
- DÖRJES, J., GADOW, S., REINECK, H.-E. & BIR SINGH, I. (1969): Die Rinnen der Jade (Südliche Nordsee). Sedimente und Makrobenthos. Senckenberg maritim 50: 5-62
- ELLIOT, M. & KINGSTON, P.F. (1987): The sublittoral benthic fauna of the estuary and Firth of Forth, Scotland. Proc Royal Soc Edinburgh, 93B: 449-465
- ESSINK, K. & KLEEF, H.L. (1988): *Marenzelleria viridis* (Verril, 1873) (Polychaeta: Spionidae): a new record from the Ems estuary (The Netherlands/Federal Republic of Germany). Zool Bijdr Leiden 38: 1-13
- ESSINK, K. & KLEEF, H.L. (1993): Distribution and life cycle of the North American spionid *Marenzelleria viridis* (Verrill, 1873) in the Ems estuary. Netherland Journal of Aquatic Ecology 27: 237-246
- ESSINK, K. & DEKKER, R. (2002): General patterns in invasion ecology tested in the Dutch Wadden Sea: the case of a brackish-marine polychaetous worm. Biol Invasions 4: 359-368
- FARKE, H. (1979): Population dynamics, reproduction and early development of *Tharyx marioni* (Polychaeta, Cirratulidae) on tidal flats of the German Bight Veröff Inst Meeresforsch Bremerhaven 18: 69-99
- GOERKE, H. (1971): Die Ernährungsweise der *Nereis*-Arten (Polychaeta, Nereidae) der deutschen Küsten. Veröff Inst Meeresforsch Bremerh 13: 1-50
- HARTMANN-SCHRÖDER, G. (1996): Annelida, Borstenwürmer, Polychaeta. In: DAHL, F. (ed.): Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise, Teil 58, 2. Aufl. Verlag G. Fischer, Jena, 648 pp
- HATTON J. & PEARCE B. (2013): The first documented record of the non-native spionid *Boccardia proboscidea* in UK waters. Mar Biodiv Rec 6, e101: 1-4, publ online
- HAUSER, B. & MICHAELIS, H. (1975): Die Makrofauna der Watten, Strände, Riffe und Wracks um den Hohen Knechtsand in der Wesermündung. Jb. Forschungsstelle Küste Norderney 1974, Bd. 26: 85-119
- JENSEN, K.T. (1992): Macrozoobenthos on an intertidal mudflat in the Danish Wadden Sea: comparisons of surveys made in the 1930s, 1940s and 1980s. Helgoländer Meeresunters. 46: 363-376
- KERCKHOF F. & FAASSE M.A. (2014): *Boccardia proboscidea* and *Boccardiella hamata* (Polychaeta: Spionidae: Polydorinae), introduced mud worms new for the North Sea and Europe, respectively. Mar Biodiv Rec 7: e76, publ online, 9 pp
- KNIGHT-JONES, P., KNIGHT-JONES, E.W., THORP, C.H. & GRAY, P.W.G. (1975): Immigrant Spirorbids (Polychaeta Sedentaria) on the Japanese *Sargassum* at Portsmouth, England. Zoologica Scripta 4: 145-149
- KÜHL, H. (1977a): *Mercierella enigmatica* (Polychaeta: Serpulidae) an der deutschen Nordseeküste. Veröff Inst Meeresforsch Bremerh 16: 99-104

- KUHLENKAMP, R. & KIND, B. (2013): Arrival of the invasive *Watersipora subtorquata* (Bryozoa) at Helgoland (Germany, North Sea) on floating macroalgae (*Himanthalia*). *Mar Biodivers Rec* 6: 1-6, published online
- MAXIMOV, A.A. (2010): Changes in bottom communities off the Eastern Gulf of Finland after Introduction of the Polychaete *Marenzelleria neglecta*. *Russ J Biol Invas* 1:11-16
- MAXIMOV, A.A. (2011): Large-Scale invasion of *Marenzelleria* spp. (Polychaeta; Spionidae) in the Eastern Gulf of Finland, Baltic Sea. *Russ J Biol Invas* 2: 11-19
- MCLUSKY, D.S., HULL, S.C. & ELLIOTT, M. (1993): Variations in the intertidal and subtidal macrofauna and sediments along a salinity gradient in the upper Forth estuary. *Neth J Aquat Ecol* 27: 101-109
- MICHAELIS, H. (1969): Makrofauna und Vegetation der Knechtsandwatten. *Jber. 1967, Forschungsstelle Norderney*, 19: 147-173
- MÜLLER, C. (2007): Konkurrenz zwischen Polychaeten im Wattenmeer. Diplomarbeit, Phillips-Universität Marburg, 101 pp
- NEIDEMAN, R., WENNGREN, J. & OLAFSSON, E. (2003): Competition between the introduced polychaete *Marenzelleria* sp. and the native amphipod *Monoporeia affinis* in Baltic soft bottoms. *Mar Ecol Prog Ser* 264: 49-55
- ORLOVA, M.I., TELESH, I.V., BEREZINA, N.A., ANTSULEVICH, A.E., MAXIMOV, A.A. & LITVINCHUK, L.F. (2006): Effects of nonindigenous species on diversity and community functioning in the eastern Gulf of Finland (Baltic Sea). *Helgol Mar Res* 60: 98-105
- PERNET, B., BARTON, M., FITZHUGH, K., HARRIS, L.H., LIZÁRRAGA, D., OHL, R. & WHITCRAFT, C.R. (2016): Establishment of the reef-forming tubeworm *Ficopomatus enigmaticus* (Fauvel, 1923) (Annelida: Serpulidae) in southern California. *Bioinvasions Rec* 5:13-19
- RADASHEVSKY, V. (2012): Spionidae (Annelida) from shallow waters around the British Islands: an identification guide for the NMBAQC Scheme with an overview of spionid morphology and biology. *Zootaxa* 3152: 1-35
- READ, G.B. (2010): Comparison and history of *Polydora websteri* and *P. haswelli* (Polychaeta: Spionidae) as mud-blister worms in New Zealand shellfish. *NZ J Mar Freshw Res* 44:83-100
- REISE, K. (1990): Historische Veränderungen in der Ökologie des Wattenmeeres. *Rheinisch-Westfälische Akad Wiss, Vorträge N 382*: 35-50
- REISE, K., GOLLASCH, S. & WOLFF, W.J. (1999): Introduced marine species of the North Sea coast. *Helgoländer Meeresunters* 52: 219-243
- SCHWINDT, E., BORTULUS, A. & IRIBARNE, O.O. (2001): Invasion of a reef-builder Polychaete: direct and indirect impacts on the native benthic community structure. *Biol Invasions* 3: 137-149
- SIKORSKI, A.V. & BICK, A. (2004): Revision of *Marenzelleria* Mesnil, 1896 (Spionidae, Polychaeta). *Sarsia* 89: 253-275
- SURUGIU, V. (2012): Systematics and ecology of species of the *Polydora*-complex (Polychaeta: Spionidae) of the Black Sea. *Zootaxa* 3518: 45-65

- WESTHEIDE, W. (1967): Monographie der Gattungen *Hesionides* Friedrich und *Microphthalmus* Mecznirow (Polychaeta, Hesionidae). Z Morph Tiere 61: 1-159
- WOLFF, W.J. (1969): Three species of *Microphthalmus* (Polychaeta) new to The Netherlands. Zool Meded, Leiden, 43: 307-311
- WOZNICZKA, A, GROMISZ, S. & WOLNOMIEJSKI, N. (2011): *Hypania invalida* (Grube, 1960), a polychaete species new for the southern Baltic estuarine area: the Szczecin Lagoon and the River Odra mouth. Aquatic Invasions 6: 39-46
- YSEBAERT, T., MEIRE, P., DE BLOCK, M., DE REGGE, N. & SOORS, J. (1997): A first record of *Marenzelleria viridis* (Verrill, 1873) (Polychaeta, Spionidae) in the Schelde Estuary (Belgium). Biol Joarb Dodonaea, 64, 1996 (1997): 176-181
- Zettler, M.L. & Röhner, M. (2004): Die Biodiversität in der deutschen Nord- und Ostsee, Bd. 3: Verbreitung und Entwicklung des Makrozoobenthos der Ostsee zwischen Fehmarnbelt und Usedom – Daten von 1839 bis 2001. BfG-1421
- Zettler, M., Daunys, D., Kotta, J. & Bick, A. (2002): History and success of an invasion into the Baltic Sea: The polychaete *Marenzelleria* cf. *viridis*, development and strategies. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): Invasive aquatic species of Europe. Distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, Boston, London, 27-52
- Zibrowius, H. & Thorp, C.H. (1989): A review of the alien serpulid and spirorbid Polychaetes in the British Isles. Cah Biol Mar 30: 271-285

ARTIKEL ZU OLIGOCHAETA (WENIGBORSTER)

- BEDDARD, F.E. (1892): A new branchiate oligochaete (*Branchiura sowerbyi*). Quarterly Journal of microscopical Science 33: 325-341 + Tafel 19
- ERSÉUS, C. & KVIST, S. (2007): COI variation in Scandinavian marine species of *Tubificoides* (Annelida: Clitellata: Tubificidae). J Mar Biol Ass U.K. 87: 1121-1126
- GEISSEN, H.-P. (1999): Bemerkungen zur Verbreitung und Ökologie des Kiemenwurms *Branchiura sowerbyi* (Oligochaeta: Tubificidae). Lauterbornia 36: 93-107
- GRABOWSKI, M. & JABŁOŃSKA, A. (2009): First records of *Branchiura sowerbyi* Beddard, 1892 (Oligochaeta: Tubificidae) in Greece. Aquatic Invasions 4: 365-367
- GRUSZKA, P. (1999): The river Odra estuary as a gateway for alien species immigration to the Baltic Sea basin. Acta hydrochim hydrobiol 27: 374–382
- HARREL, R.C. (2004): Systematic and ecological notes on *Tubificoides heterochaetus* (Oligochaeta: Tubificidae) from the Neches River estuary, Texas. Texas J Sci 56: 263-267
- MANN, K.H. (1958): Occurrence of an exotic oligochaete *Branchiura sowerbyi* Beddard, 1892, in the River Thames. Nature 182, 732

- MILBRINK, G. & TIMM, T. (2001): Distribution and dispersal capacity of the Ponto-Caspian tubificid oligochaete *Potamotrix moldaviensis* Vejdovsky et Mrázek, 1903 in the Baltic Sea Region. *Hydrobiologia* 463: 93-102
- SCHUCHARDT, B., HAESLOOP, U. & SCHIRMER, M. (1993): The tidal freshwater reach of the Weser estuary: Riverine or estuarine? *Neth J Aquat Ecol* 27: 215-226
- SEYS, J., VINCX, M. & MEIRE, P. (1999): Spatial distribution of oligochaetes (Clitellata) in the tidal freshwater and brackish parts of the Schelde estuary (Belgium). *Hydrobiologia* 406: 119-132
- TOBIAS, W. (1972): Ist der Schlammröhrenwurm *Branchiura sowerbyi* Beddard 1892 (Oligochaeta: Tubificidae) ein tropischer Einwanderer im Untermain? *Nat Mus* 102: 93-107
- WOŹNICZKA A., GROMISZ S., WOLNOMIEJSKI N. (2011): *Hypania invalida* (Grube, 1960), a polchaet species new for the southern Baltic estuarine area: the Szczecin Lagoon and the River Odra mouth. *Aquatic Invasions* 6: 39-46
- ZETTLER, M. (1996): Erstnachweis von *Branchiura sowerbyi* Beddard, 1892 (Oligochaeta: Tubificidae) in Mecklenburg-Vorpommern. *Lauterbornia* 26, 99-101

ÜBERSICHTSARTIKEL UND LISTEN MIT ANGABEN ZU POLYCHAETA UND OLIGOCHAETA

- ENO, N.C., CLARK, R.A. & SANDERSON, W.G. (eds.) (1997): Non-native marine species in British waters: a review and directory. Joint Nature Conservation Committee, Peterborough, 152 pp
- GERLACH, S.A. (2000): Die Biodiversität in der deutschen Nord- und Ostsee, Bd. 1, Checkliste der Fauna der Kieler Bucht und eine Bibliographie zur Biologie und Ökologie der Kieler Bucht. Bundesanstalt für Gewässerkunde, Koblenz, 376 pp
- GITTENBERGER, A., RENSING, M., STEGENGA, H. & HOEKSEMA, B.W. (2010): Native and non-native species of hard substrata in the Dutch Wadden Sea. *Ned Faun Meded* 33: 21-76
- GOLLASCH, S. & NEHRING, S. (2006): National checklist for aquatic species in Germany *Aquatic invasions* 1: 245–269
- GOLLASCH, S., HAYDAR, D., MINCHIN, D., WOLFF, W.J. & REISE, K. (2009): Introduced Aquatic Species of the North Sea Coasts and Adjacent Brackish Waters. In: RILOV, G. & CROOKS, J.A. (eds.): *Biological invasions in marine ecosystems*, Chapter 29: 507-528 *Ecological Studies* 204, Springer-Verlag Berlin Heidelberg 2008
- HAYWARD, P.J. & RYLAND, J.S. (1990): *The Marine fauna of the British Isles and North-West Europe*, Vol. 1 & 2. Oxford Science Publications, Clarendon Press, Oxford, 1990
- JENSEN, K. & KNUDSEN, J. (2005): A summary of alien marine benthic invertebrates in Danish waters. *Oceanol Hydrobiol Stud* 34, Suppl 1: 137-162
- KERCKHOF, F., HAELTERS, J. & GOLLASCH, S. (2007): Alien species in the marine and brackish water ecosystems: the situation in Belgian waters. *Aquatic Invasions* 2: 243-257

- NEHRING, S. (2006): Four arguments why so many alien species settle into the estuaries, with special reference to the German river Elbe. *Helgol Mar Res* 60: 127-134
- NEHRING, S. & LEUCHS, H. (1999a): Neozoa (Makrobenthos) an der deutschen Nordseeküste. Eine Übersicht. Bundesanstalt für Gewässerkunde, Koblenz. 131 pp
- NEHRING, S., REISE, K., DANKERS, N. & KRISTENSEN, P.S. (2009): Alien species. Thematic Report No. 7. In: MARENCIC, H. & VLAS, J. DE (Eds), 2009. Quality Status Report 2009. Wadden Sea Ecosystem No. 25. Common Wadden Sea Secretariat, Trilateral Monitoring and Assessment Group, Wilhelmshaven, Germany
- POST, D. & LANDMANN, M. (1994): Verbreitungsatlas der Fließgewässerfauna in Ostfriesland. Staatl Amt Wasser Abfall, 141 S.
- TITTIZER, T., SCHÖLL, F., BANNING, M., HAYBACH, A. & SCHLEUTER, M. (2000): Aquatische Neozoen im Makrobenthos der Binnenwasserstraßen Deutschlands. *Lauterbornia* 39: 1-72
- WOLFF, W.J. (2005): Non-indigenous marine and estuarine species in The Netherlands. *Zool Meded* 79: 1-116
- Zettler, M.L. & Röhner, M. (2004): Die Biodiversität in der deutschen Nord- und Ostsee, Bd. 3: Verbreitung und Entwicklung des Makrozoobenthos der Ostsee zwischen Fehmarnbelt und Usedom – Daten von 1839 bis 2001. BfG-1421

Listen und Internetquellen zu Annelida

www.frammandearter.se
www.nobanis.org