

LITERATUR

CRUSTACEA (KREBSE)

ARTIKEL ZU CLADOCERA (WASSERFLÖHE)

- ANTSULEVICH, A. & VÄLIPAKKA, P. (2000): New important food object of the Baltic herring in the Gulf of Finland. *Internat Rev Hydrobiol* 85: 605-615
- BIELECKA, L., ZMIJEWSKA, M. I. & SZYMBORSKA, A. (2000): A new predatory cladoceran *Cercopagis (Cercopagis) pengoi* (Ostroumov 1891) in the Gulf of Gdansk. *Oceanologia* 42: 371-374
- BIRNBAUM, C. (2006): NOBANIS – Invasive Alien Species Fact Sheet – *Cercopagis pengoi*. – From: Online Database of the European Network on Invasive Alien Species – NOBANIS www.nobanis.org (04.06.2013)
- GOROKHOVA, E., ALADIN, N. & DUMONT, H.J. (2000): Further expansion of the genus *Cercopagis* (Crustacea, Branchiopoda, Onychiopoda) in the Baltic Sea, with notes on the taxa present and their ecology. *Hydrobiologia* 429: 207-218
- JAŹDŹEWSKI, K. & KONOPACKA, A. (2002): Invasive Ponto-Caspian species in waters of the Vistula and Oder Basins and the Southern Baltic Sea. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, Boston, London, 384-398
- KRYLOV, P.I., BYCHENKOV, D.E., PANOV, V.E., RODIONOVA, N.V. & TELESH, I.V. (1999): Distribution and seasonal dynamics of the Ponto-Caspian invader *Cercopagis pengoi* (Crustacea, Cladocera) in the Neva Estuary (Gulf of Finland). *Hydrobiologia* 393: 227-232
- OJAVEER, H. (1997): *Cercopagis pengoi*. In: *Baltic Sea Alien Species Database*, <http://www.corpi.ku.lt/nemo/mainnemo.html>
- OJAVEER, H. & LUMBERG, A. (1995): On the role of *Cercopagis (Cercopagis) pengoi* (Ostroumov) in the Pärnu Bay and the NE part of the Gulf of Riga ecosystem. *Proc Estonian Acad Sci Ecol* 5: 20-25
- ORLOVA, M.I., TELESH, I.V., BEREZINA, N.A., ANTSULEVICH, A.E., MAXIMOV, A.A. & LITVINCHUK, L.F. (2006): Effects of nonindigenous species on diversity and community functioning in the eastern Gulf of Finland (Baltic Sea). *Helgol Mar Res* 60: 98-105
- TELESH, I.V. & OJAVEER, H. (2002): The predatory water flea *Cercopagis pengoi* in the Baltic Sea: Invasion history, distribution and implications to ecosystem dynamics. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, Boston, London, 62-65

ARTIKEL ZU COPEPODA (RUDERFÜßER)

- AGUIRRE-MACEDO, M.L. & KENNEDY, C.R. (1999): Diversity of metazoan parasites of the introduced oyster species *Crassostrea gigas* in the Exe Estuary. *J Mar Biol Ass UK* 79: 57-63
- ARNDT, H. & SCHNESE, W. (1986): Population dynamics and production of *Acartia tonsa* (Copepoda: Calanoida) in the Darss-Zingst estuary, southern Baltic. *Ophelia*, Suppl 4: 329-334
- BAIRD, R.H., BOLSTER, G.C. & COLE, H.A. (1951): *Mytilicola intestinalis*, Steuer, in the European Flat Oyster (*Ostrea edulis*). *Nature* 168: 560
- BARETTA, J. W. & MALSCHAERT, J. F. P. (1988): Distribution and abundance of the zooplankton of the Ems estuary (North Sea). *Neth J Sea Res* 22: 69-81
- BEHREND, G., KORSHENKO, A. & VIITASALO, M. (1997): Morphological aberrations in females of the genus *Acartia* (Copepoda, Calanoida) in the Baltic Sea. *Crustaceana* 70: 596 ff.
- BĒRZINŠ, B. (1940): *Acartia tonsa* in the Gulf of Riga. *Folia Zoologica et Hydrobiologica* 10: 484-486
- BRYLINSKI, J.-M. (1981): Report on the presence of *Acartia tonsa* Dana (Copepoda) in the harbour of Dunkirk (France) and its geographical distribution in Europe. *J Plankton Res* 3: 255-260
- BRYLINSKI, J.-M., ANTAJAN, E., RAUD, T. & VINCENT, D. (2012): First record of the Asian copepod *Pseudodiaptomus marinus* Sato, 1913 (Copepoda: Calanoida: Pseudodiaptomidae) in the southern bight of the North Sea along the coast of France. *Aquatic Invasions* 7: 577-584
- CASPERS, H. (1939): Über Vorkommen und Metamorphose von *Mytilicola intestinalis* Steuer (Copepoda paras.) in der südlichen Nordsee. *Zool Anz* 126: 161-171
- CLANZIG, S. (1989): Invertébrés d'introduction récente dans les lagunes méditerranéennes du Languedoc-Roussillon (France). *Bull. Soc. Zool. France* 114: 151-152
- CORNILS A., WEND-HECKMANN B. (2015): First report of the planktonic copepod *Oithona davisae* in the northern Wadden Sea (North Sea): Evidence for recent invasion? *Helgol Mar Res* 69 (2): 243-248
- DAVEY, J.T. & GEE, J.M. (1976): The occurrence of *Mytilicola intestinalis* Steuer, an intestinal copepod parasite of *Mytilus*, in the south-west of England. *J mar biol Ass U.K.* 56: 85-94
- DETHLEFSEN, V. (1972): Zur Parasitologie der Miesmuschel (*Mytilus edulis* L., 1758). *Ber dt wiss Kommn Meeresforsch* 22: 344-371
- DETHLEFSEN, V. (1975): The influence of *Mytilicola intestinalis* Steuer on the meat content of the mussel *Mytilus edulis* L. *Aquaculture* 6: 83-97
- ELLENBY, C. (1947): A copepod parasite of the mussel new to the British fauna. *Nature* 159: 645-646
- ELSNER, N.O. (2009): Eingeschleppte Copepoden der Gattung *Mytilicola* im Darm von Miesmuscheln und Austern im Seegebiet der Nordseeinsel Sylt. Diplomarbeit, Univ. Marburg, 65 pp

- ELSNER, N.O., JACOBSEN, S., THIELTGES, D.W. & REISE K. (2011): Alien parasitic copepods in mussels and oysters of the Wadden Sea. *Helgol Mar Res* 65: 299-307
- FLEMINGER, A. & HENDRIX KRAMER, S. (1988): Recent introduction of an Asian copepod, *Pseudodiaptomus marinus* (Copepoda: Calanoida), into southern California embayments. *Mar Biol* 98: 535-541
- GEE, J.M. & DAVEY, J.T. (1986): Stages in the life history of *Mytilicola intestinalis* Steuer, a copepod parasite of *Mytilus edulis* (L.), and the effect of temperature on their rates of development. *J Cons Int Explor Mer* 42: 254-264
- HOLMES, J.M.C. & MINCHIN, D. (1995): Two exotic copepods imported into Ireland with the Pacific Oyster *Crassostrea gigas* (Thunberg). *Ir Nat J* 25: 17-19
- JENSEN, K.R. (2010b): NOBANIS – Invasive Alien Species Fact Sheet – *Mytilicola intestinalis*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (06.06.2013)
- JHA, U., JETTER, A., LINDLEY, J.A., POSTEL, L. & WOOTTON, M. (2013): Extension of distribution of *Pseudodiaptomus marinus*, an introduced copepod, in the North Sea. *Mar Biodivers Rec* 6, e53, 3 pp, published online
- KLIE, W. (1930): Neues zur Crustaceen-Fauna Nordwestdeutschlands. *Abh Nat Ver Bremen* 28: 271-276
- KORRINGA, P. (1968): On the ecology and distribution of the parasitic copepod *Mytilicola intestinalis* Steuer. *Bijdr dierkd* 38: 47-57
- LAUCKNER, G. (1983): Diseases of Mollusca, Bivalvia (Copepoda, p. 817-829). In: KINNE, O.: *Diseases of Marine Animals*, Vol. 2, 1083 pp
- MEYER, P.-F. & MANN, H. (1950): Beiträge zur Epidemiologie und Physiologie des parasitischen Copepoden *Mytilicola intestinalis*. *Arch Fischereiwiss* 2: 120-134
- MEYER-WAARDEN, P.-F. & Mann, H. (1954): Der Befall von *Mytilus edulis* durch *Mytilicola intestinalis* in den deutschen Wattengebieten 1950-1953. *Ber Dtsch Wiss Komm Meeresforsch* 13: 347-362
- OLAZABAL, A. DE & TIRELLI, V. (2011): First record of the egg-carrying calanoid copepod *Pseudodiaptomus marinus* in the Adriatic Sea. *Mar Biodivers Rec* 4, e85:1-4
- POGODA B., JUNGBLUT S., BUCK B.H., HAGEN W. (2012): Infestation of oysters and mussels by mytilicolid copepods: differences between natural coastal habitats and two offshore cultivation sites in the German Bight. *J Appl Ichtyol* 28: 756-765
- REMY, P. (1927): Note sur un Copépode de l'eau saumâtre du Canal de Caen à la mer. *Ann biol lacustre* 15: 169-186
- SCHEIBEL, W. (1973): Quantitativ-ökologische Untersuchungen am uferfernen Mesopsammon in der Kieler Bucht. *Kieler Meeresforsch* 29: 58-68
- SCHEIBEL, W. (1974): *Ameira divagans* Nicholls, 1939 (Copepoda Harpacticoidea). *Mikrofauna Meeresboden* 38: 1-10
- SCHEIBEL, W. & RUMOHR, H. (1979): Meiofaunaentwicklung auf künstlichen Weichböden in der Kieler Bucht. *Helgoländer Meeresunters* 32: 305-312

- SEI, S. & FERRARI, I. (2008): First report of the occurrence of *Acartia tonsa* (Copepoda: Calanoida) in the Lesina lagoon (south Adriatic Sea – Mediterranean Sea). *Mar Biodivers Rec* 1; e37; 2 pages; published online
- STEELE, S. & MULCAHY, M.F. (2001): Impact of the copepod *Mytilicola orientalis* on the Pacific oyster *Crassostrea gigas* in Ireland. *Dis Aquat Org* 47: 145-149
- STEUER, A. (1902): *Mytilicola intestinalis* n. gen. n. sp. aus dem Darm von *Mytilus galloprovincialis* Lam. *Zool Anz* 25: 635-637
- STOCK, J.H. (1993): Copepoda (Crustacea) associated with commercial and non-commercial Bivalvia in the East Scheldt, The Netherlands. *Bijdr Dierkunde* 63: 61-64
- THEISEN, B.F. (1966): *Mytilicola intestinalis* Steuer in Danish Waters 1964-65. *Medd fra Danmarks Fiskeri- og Havundersoegelser* 4: 327-337
- THEISEN, B.F. (1987): *Mytilicola intestinalis* Steuer and the condition of its host *Mytilus edulis*. *Ophelia* 27: 77-86
- THIELTGES, D.W., KRAKAU, M., ANDRESEN, H., FOTTNER, S. & REISE, K. (2006): Macroparasite community in molluscs of a tidal basin in the Wadden Sea. *Helgol Mar Res* 60: 307-316
- TISELIUS, P., BORG, C.M.A., HANSEN, B.W., HANSEN, P.J., NIELSEN, T.G. & VISMANN, B. (2008): High reproduction, but low biomass: mortality estimates of the copepod *Acartia tonsa* in a hyper-eutrophic estuary. *Aquat Biol* 2: 93-103
- ZAIKO, A. (2004b): *Acartia tonsa*. In: Baltic Sea Alien Species Database. S. Olenin, E. Leppakoski and D. Dauny (eds.). www.corpi.ku.lt/nemo/alien_species_directory.html

ARTIKEL ZU CIRRIPEDIA (RANKENFÜßER)

- BARNES, H. & BARNES, M. (1960): Recent spread and present distribution of the barnacle *Elminius modestus* Darwin in North-West Europe. *Proc Zool Soc London* 135: 137-145
- BISHOP, M.W.H. (1947): Establishment of an immigrant barnacle in British coastal waters *Nature* 159: 501-502
- BISHOP, M.W.H. & CRISP, D.J. (1958): The distribution of the barnacle *Elminius modestus* Darwin in France. *Proc Zool Soc London* 131: 109-134
- BUCKERIDGE, J.S. & NEWMAN, W.A. (2010): A review of the subfamily Eliminiinae (Cirripedia: Thoracica: Austrobalanidae), including a new genus, *Protelminius* nov., from the Oligocene of New Zealand. *Zootaxa* 2349: 39-54
- CRISP, D.J. (1958): The spread of *Elminius modestus* Darwin in North-West Europe. *J mar biol Ass U.K.* 37: 483-520
- DEN HARTOG, C. (1953): Immigration, dissemination and ecology of *Elminius modestus* Darwin in the North Sea, especially along the Dutch coast. *Beaufortia* 4: 1-20

- HARMS, J. (1999): The neozoan *Elminius modestus* Darwin (Crustacea, Cirripedia): Possible explanations for its successful invasion in European water. Helgoländer Meeresunters 52: 337-345
- JENSEN, K.R. (2010c): NOBANIS – Invasive Alien Species Fact Sheet – *Elminius modestus*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (10.06.2013)
- JENSEN, K.R. (2010d): NOBANIS – Invasive Alien Species Fact Sheet – *Balanus improvisus*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (13.06.2013)
- KÜHL, H. (1954): Über das Auftreten von *Elminius modestus* Darwin in der Elbmündung. Helgoländer Meeresunters 5: 53-56
- KÜHL, H. (1963): Die Verbreitung von *Elminius modestus* Darwin (Cirripedia Thoracica) an der deutschen Küste. Crustaceana 5: 99-111
- LAWSON, J., DAVENPORT, J. & WHITAKER, A. (2004): Barnacle distribution in Lough Hyne Marine Nature Reserve: a new baseline and an account of invasion by the introduced Australasian species *Elminius modestus* Darwin. Estuar Coast Shelf Sci 60: 729-735
- LEPPÄKOSKI, E. (1999): *Balanus improvisus* (Darwin 1854). In: GOLLASCH, S., MINCHIN, D., ROSENTHAL, H. & VOIGT, M. (Hrsg.): Exotics across the ocean. Case histories on introduced species: their general biology, distribution, range expansion and impact. Kiel 1999, 49-54
- LUTHER, G. (1987): Seepocken der deutschen Küstengewässer. Helgoländer Meeresunters 41: 1- 43
- OLSZEWSKA, A. (2000): *Mya arenaria* L., a new and unusual substratum for *Balanus improvisus* Darwin. Oceanologia 42: 119-121
- O'RIORDAN, R.M., CULLOTY, S., DAVENPORT, J. & MCALLEN, R. (2009): Increases in the abundance of the invasive barnacle *Austrominius modestus* on the Isle of Cumbrae, Scotland. Mar Biodivers Rec 2: 1-4
- STUBBINGS, H.G. (1950): Earlier records of *Elminius modestus* Darwin in British waters. Nature 166: 277-278
- WITTE, S., BUSCHBAUM, C., VAN BEUSEKOM, J.E.E. & REISE, K. (2010): Does climatic warming explain why an introduced barnacle finally takes over after a lag of more than 50 years? Biol Invasions 12: 3579-3589
- ZAICO, A. (2005b): *Balanus improvisus*. In: Baltic Sea Alien Species Database. S. Olenin, E. Leppakoski and D. Dauny (eds.). www.corpi.ku.lt/nemo/alien_species_directory.html

ARTIKEL TANAIIDACEA (SCHERENASSELN)

- BAMBER R.N. (2014): Two new species of *Sinelobus* Sieg, 1980 (Crustacea: Tanaidacea: Tanaididae), and a correction to the higher taxonomic nomenclature. J Nat Hist 48: 2049-2068

VAN HAAREN, T. & SOORS, J. (2009): *Sinelobus stanfordi* (Richardson, 1901): A new crustacean invader in Europe. *Aquatic Invasions* 4: 703-711

ARTIKEL ZU MYSIDACEA (SCHWEBGARNELEN)

ARBACIAUSKAS, K. (2002): Ponto-Caspian amphipods and mysids in the inland waters of Lithuania: History of introduction, current distribution and relations with native Malacostracans. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, Boston, London, 104-115

GERGS, R., HANSELMANN, A.J., EISELE, I. & ROTHHAUPT, K.-O. (2008): Autecology of *Limnomysis benedeni* Czerniavsky, 1882 (Crustacea: Mysida) in Lake Constance, Southwestern Germany. *Limnologia* 38: 139-146

HAESLOOP, U. (2001): Einige bemerkenswerte Makrovertebraten-Funde aus Gewässern des Großraumes Bremen. *Lauterbornia* 41: 55-59

JANAS, U. & WYSOCKI, P. (2005): *Hemomysis anomala* G.O. Sars, 1907 (Crustacea, Mysidacea) – first record in the Gulf of Gdansk. *Oceanologia* 47: 405-408

JENSEN, K.R. (2010e): NOBANIS – Invasive Alien Species Fact Sheet – *Hemimysis anomala*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (17.06.2013)

KETELAARS, H.A.M., LAMBREGTS-VAN DE CLUNDERT, F.E., CARPENTIER, C.J., WAGENVOORT, A.J. & HOOGENBOEZEM, W. (1999): Ecological effects of the mass occurrence of the Ponto-Caspian invader, *Hemimysis anomala* G.O.Sars, 1907 (Crustacea: Mysidacea), in a freshwater storage reservoir in the Netherlands, with notes on its autecology and new records. *Hydrobiologia* 394: 233-248

MICHELS, U. (2005): *Limnomysis benedeni* (Crustacea: Mysidacea) neu für die Untere Oder in Polen. *Lauterbornia* 55: 83-87

OVČARENKO, I., AUDZIJONYTE, A. & GASIŪNAITE, Z.R. (2006): Tolerance of *Paramysis lacustris* and *Limnomysis benedeni* (Crustacea, Mysida) to sudden salinity changes: implications for ballast water treatment. *Oceanologia* 48: 231-242

ROTH, H. & ZETTLER M.L. (2015): Morphologische und ökologische Eigenschaften allochtoner Mysida aus der Pontokaspis. *Lauterbornia* 80: 51-68

RUDOLPH, K. & ZETTLER, M.L. (2003): Erste Nachweise der Schwebgarnele *Hemimysis anomala* Sars, 1907 (Crustacea, Mysidacea) in Wasserstrassen im Nordosten Deutschlands. *Sitzungsberichte der Gesellschaft naturforschender Freunde zu Berlin* 42: 78-83

VERSLYCKE, T., JANSSEN, C., LOCK, K. & MEES, J. (2000): First occurrence of the Pontocaspian invader *Hemimysis anomala* (Sars, 1907) in Belgium (Crustacea: Mysidacea). *Belg J Zool* 130: 157-158

WITTMANN, K.J. (2011): Datasheet for CABI Invasive species compendium, invasive aquatic

species – *Limnomysis benedeni*

www.meduniwien.ac.at/user/karl.wittmann/docs/2011/wittmann_2011.pdf

(21.10.2011)

ZETTLER M. (2015): Kurze Notiz über die Ankunft von *Echinogammarus trichiatus* im Ostseegebiet und den Erstrnachweis von *Paramysis lacustris* in Deutschland. *Lauterbornia* 79: 151-156

Zettler, M.L. & Daunys, D. (2007): Long-term macrozoobenthos changes in a shallow boreal lagoon: Comparison of a recent biodiversity inventory with historical data. *Limnologica* 37: 170-185

ARTIKEL ZU ISOPODA (ASSELN)

CHAPMAN J.W. & CARLTON J.T. (1994): Predicted discoveries of the introduced isopod *Synidothea laevidorsalis* (Miers, 1881). *J Crust Biol* 14: 700-714

NEHRING, S. & LEUCHS, H. (2000): Neozoen im Makrozoobenthos der Brackgewässer an der deutschen Nordseeküste. *Lauterbornia* 39: 73-116

POORE G.C.B. (1996): Species differentiation in *Synidothea* (Isopoda: Idotheidae) and recognition of introduced marine species: a reply to Chapman and Carlton. *J Crust Biol* 16: 384-394

SOORS, J., FAASSE, M., STEVENS, M., VERBESSEM, I., DE REGGE, N. & VAN DEN BERGH, E. (2010): New crustacean invaders in the Schelde estuary (Belgium). *Belg J Zool* 140: 3-10

WITTFOTH, A.K.J. & ZETTLER, M.L. (2013): The application of a Biopollution Index in German Baltic estuarine and lagoon waters. *Management of Biol Invasions* 4: 43-50

ZETTLER, M.L. (1998): Zur Verbreitung der Malacostraca (Crustacea) in den Binnen- und Küstengewässern von Mecklenburg-Vorpommern. *Lauterbornia* 32: 49-65

ARTIKEL ZU AMPHIPODA (FLOHKREBSE)

ALMÓN, B., PÉREZ, J., BANÓN, R. & TRIGO, J. (2014): First record of *Caprella mutica* from the Iberian Peninsula: expansion southwards in European waters. *Marine Biodiversity Records* 7: 1-4

ASHTON, G.V., STEVENS, M.I., HART, M.C., GREEN, D.H., BURROWS, M.T., COOK, E.J. & WILLIS, K.J. (2008): Mitochondrial DNA reveals multiple Northern Hemisphere introductions of *Caprella mutica* (Crustacea, Amphipoda). *Mol Ecol* 17: 1293-1303

BAÇELA, K. & KONOPACKA, A. (2005): The life history of *Pontogammarus robustoides*, an alien amphipod species in Polish waters. *Journal of Crustacean Biology* 25: 190-195

- BAĆELA, K., GRABOWSKI, M. & KONOPACKA, A. (2008): *Dikerogammarus villosus* (Sowinski, 1894) (Crustacea, Amphipoda) enters Vistula – the biggest river in the Baltic basin. *Aquatic Invasions* 3: 95-98
- BAĆELA, K., KONOPACKA, A. & GRABOWSKI, M. (2009): Reproductive biology of *Dikero-gammarus haemobaphes*: an invasive gammarid (Crustacea: Amphipoda) colonizing running waters in Central Europe. *Biol Invasions* 11: 2055-2066
- BEERMANN, J. & FRANKE, H.-D. (2012): Differences in resource utilization and behaviour between coexisting *Jassa* species (Crustacea, Amphipoda). *Mar Biol* 159: 951-957
- BEREZINA, N. A. (2007): Expansion of the North American amphipod *Gammarus tigrinus* Sexton, 1939 to the Neva Estuary (easternmost Baltic Sea). *Oceanologia* 49: 129-135
- BIJ DE VAATE, A., JAZDZEWSKI, K., KETELAARS, H.A.M., GOLLASCH, S. & VAN DER VELDE, G. (2002): Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in Europe. *Can J Fish Aquat* 59: 1159-1174
- BOCK, K.-D. (1967): Experimente zur Ökologie von *Orchestia platensis* Kröyer *Z Morph Ökol Tiere* 58: 405-428
- BOETS, P., LOCK, K., TEMPELMAN D., VAN HAAREN T.V., PLATVOET D., GOETHALS P.L.M. (2012): First occurrence of the Ponto-Caspian amphipod *Echinogammarus trichiatus* (Martynov, 1932) (Crustacea: Gammaridae) in Belgium. *BiolInvasion Rec* 1: 115-120
- BOETS, P., LOCK, K. & GOETHALS, L.M. (2012): Assessing the importance of alien macro-crustacea (Malacostraca) within macroinvertebrate assemblages in Belgian coastal harbours. *Helgol Mar Res* 66: 175-187
- BULNHEIM, H.-P. (1976): *Gammarus tigrinus*, ein neues Faunenelement der Ostseeförde Schlei. *Schr Naturw Ver Schlesw-Holst* 46: 79-84
- BUSCHBAUM, C. & GUTOW, L. (2005): Mass occurrence of an introduced crustacean (*Caprella* cf. *mutica*) in the south-eastern North Sea. *Helgol Mar Res* 59: 252-253
- CONLAN, K.E. (1990): Revision of the crustacean amphipod genus *Jassa* Leach (Corophioidea: Ischyroceridae). *Can J Zool* 68: 2031-2075
- COOK, E.J., JAHNKE, M., KERCKHOF, F., MINCHIN, D., FAASSE, M., BOOS, K. & ASHTON, G. (2007): European expansion of the introduced amphipod *Caprella mutica* Schurin 1935. *Aquatic Invasions* 2: 411-421
- CURRY, A., GRAYSON, R.F. & MILLIGAN, T.D. (1972): New British records of the semi-terrestrial amphipod *Orchestia cavimana*. *Freshwat Biol* 2: 55-56
- CRAWFORD, G.I. (1935): *Corophium curvispinum*, G.O. Sars, var. *devium*, Wundsch, in England. *Nature* 136: 685-686
- DANELIYA, M.E. & LAAKKONEN H. (2012): The Japanese skeleton shrimp *Caprella mutica* (Amphipoda: Caprellidae) in Sweden (Eastern Skagerrak). *Mar Biodivers Rec* 5; e36; 6 pages <http://dx.doi.org/10.1017/S1755267212000243>, published online: 25. April 2012

- DAUNYS, D. & ZETTLER, M. L. (2006): Invasion of the North American amphipod (*Gammarus tigrinus* Sexton, 1939) into the Curonian Lagoon, South Eastern Baltic Sea. *Acta Zoologica Lituanica* 16: 20-26
- DEN HARTOG, C. (1963): The amphipods of the Deltaic region of the rivers Rhine, Meuse and Scheldt in relation to the hydrography of the area, Part II. The Talitridae. *Neth J Sea Res* 2: 40-67
- FAASSE, M. & VAN MOORSEL, G. (2000): Nieuwe en minder bekende Vlokreftjes van sublitorale harde bodems in het Deltagebied (Crustacea: Amphipoda: Gammaridea). *Ned Faun Meded* 11: 19-44
- FAASSE, M. & VAN MOORSEL, G. (2003): The North-American amphipod, *Melita nitida* Smith, 1873 and *Incisocalliope aestuarius* (Watling and Maurer, 1973) (Crustacea: Amphipoda: Gammaridea), introduced to the Western Scheldt estuary (The Netherlands). *Aquatic Ecology* 37: 13-22
- EGGERS, T.O. & ANLAUF, A. (2005): *Obesogammarus crassus* (G.O.Sars, 1894) (Crustacea: Amphipoda) erreicht die Elbe. *Lauterbornia* 55: 125-128
- FLECKENSTEIN, N. (2011): The role of the immigrant *Caprella mutica* in coastal ecosystems of the German Bight. Diploma Thesis, C.v.O. University Oldenburg, 28 pp
- GRABOWSKI, M. (2011): NOBANIS – Invasive Alien Species Fact Sheet – *Pontogammarus robustoides*. – From: Online Database of the European Network on Invasive Alien Species - NOBANIS www.nobanis.org (08.07.2013)
- GRABOWSKI, M., KONOPACKA, A., JAZDZEWSKI, K. & JANOWSKA, E. (2006): Invasion of alien gammarid species and retreat of natives in the Vistula Lagoon (Baltic Sea, Poland). *Helgol Mar Res.* 60: 90-97
- GRABOWSKI, M., JAŹDŹEWSKI, K. & KONOPACKA, A. (2007a): Alien Crustacea in Polish waters - Amphipoda. *Aquatic Invasions* 2: 25-38
- GRABOWSKI, M., BACELA, K. & KONOPACKA, A. (2007b): How to be an invasive gammarid (Amphipoda: Gammaroidea) – comparison of life history traits. *Hydrobiologia* 590: 75-84
- GRUSZKA, P. (1999): The river Odra estuary as a gateway for alien species immigration to the Baltic Sea basin. *Acta hydrochim hydrobiol* 27: 374–382
- GRUSZKA, P. & WOŹNICZKA, A. (2008): *Dikerogammarus villosus* (Sowinski, 1894) in the River Odra estuary – another invader threatening Baltic Sea coastal lagoons. *Aquatic Invasions* 3: 395-403
- GUMULIAUSKAITĖ, S. & ARBAČIAUSKAS, K. (2008): The impact of the invasive Ponto-Caspian amphipod *Pontogammarus robustoides* on littoral communities in Lithuanian lakes. *Hydrobiologia* 599: 127-134
- HERKÜL, K., KOTTA, J. & KOTTA, I. (2006): Distribution and population characteristics of the alien talitrid amphipod *Orchestia cavimana* in relation to environmental conditions in the Northeastern Baltic Sea. *Helgol Mar Res.* 60: 121-126
- HERKÜL, K., KOTTA, J., PÜSS, T. & KOTTA, I. (2009): Crustacean invasion in the Estonian coastal sea. *Estonian Journal of Ecology* 58 (4): 313-323

- HEROLD, W. (1925): Der Amphipode *Orchestia cavimana* Heller in Pommern. Abh u Ber Pomm Naturf Ges 6: 109-110
- HYNES, H.B.N. (1955): Distribution of some freshwater Amphipoda in Britain. Verh internat Ver theor angew Limnol 12: 620-628
- JAŹDŹEWSKI, K., KONOPACKA, A. & GRABOWSKI, M. (2005): Native and alien malacostracan Crustacea along the Polish Baltic Sea coast in the twentieth century. Oceanol Hydrobiol Stud 34, Suppl 1: 175-193
- JENSEN, K.R. (2010f): NOBANIS – Invasive Alien Species Fact Sheet – *Gammarus tigrinus*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (02.07.2013)
- JENSEN, K.R. (2010g): NOBANIS – Invasive Alien Species Fact Sheet – *Corophium sextonae*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (04.07.2013)
- JENSEN, K.R. (2010h): NOBANIS – Invasive Alien Species Fact Sheet – *Platorchestia platensis*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (08.07.2013)
- KALINKINA, N.M. & BEREZINA, N.A. (2010): First record of *Pontogammarus robustoides* Sars, 1894 (Crustacea: Amphipoda) in the Gulf of Riga (Baltic Sea). Aquatic Invasions 5, Suppl 1: 5-7
- KINZELBACH, R. (1972): Zur Verbreitung und Ökologie des Süßwasser-Strandfloh *Orchestia cavimana* Heller, 1865 (Crustacea: Amphipoda: Talitridae). Bonner Zoologische Beiträge 3 (23): 267-282
- KLEIN, G. (1969): Amphipoden aus der Wesermündung und der Helgoländer Bucht, mit Beschreibung von *Talorchestia frisiae* n.sp. Veröff Inst Meeresforsch Bremerh 11: 173-194
- KONOPACKA, A. (2003): Further step to the west – *Obesogammarus crassus* (G.O.Sars, 1894) (Crustacea, Amphipoda) already in the Szczecin Lagoon. Lauterbornia 48: 67-72
- KONOPACKA, A., GRABOWSKI, M., BACELA-SPYCHALSKA, K. & REWICZ, T. (2009): *Orchestia cavimana* Heller, 1865 (Amphipoda: Talitridae) enters freshwater inland habitats in the Vistula River, Poland. Aquatic Invasions 4: 689-691
- KOTTA, J., PÄRNOJA, M., KATAJISTO, T., LEHTINIEMI, M., MALAVIN, S.A., REISALU, G., PANOV, V.E. (2013): Is a rapid expansion of the invasive amphipod *Gammarus tigrinus* Sexton, associated with its niche selection: a case study in the Gulf of Finland, the Baltic Sea. Aquatic Invasions 8: 319-332
- MARTENS, A., EGGERS, T.O. & GRABOW, K. (1999): Erste Funde von *Pontogammarus robustoides* (SARS) im Mittellandkanal (Crustacea: Amphipoda). Lauterbornia 35: 39-42
- MÜLLER, O., ZETTLER, M.L. & GRUSZKA, P. (2001): Verbreitung und Status von *Dikerogammarus villosus* (Sovinski 1894) (Crustacea: Amphipoda) in der mittleren und unteren Strom-Oder und den angrenzenden Wasserstraßen. Lauterbornia 41: 105-112

- NEUHAUS, E. (1933): Studien über das Stettiner Haff und seine Nebengewässer. Zeitschr f Fischerei u deren Hilfswiss 31: 427-489
- ORAV-KOTTA, H., KOTTA, J., HERKÜL, K., KOTTA, I. & PAALME, T. (2009): Seasonal variability in the grazing potential of the invasive amphipod *Gammarus tigrinus* and the native amphipod *Gammarus salinus* (Amphipoda: Crustacea) in the northern Baltic Sea. Biol Invasions 11: 597-608
- ORLOVA, M.I., TELESH, I.V., BEREZINA, N.A., ANTSULEVICH, A.E., MAXIMOV, A.A. & LITVINCHUK, L.F. (2006): Effects of nonindigenous species on diversity and community functioning in the eastern Gulf of Finland (Baltic Sea). Helgol Mar Res 60: 98-105
- PACKALÉN, A., KORPINEN, S. & LEHTONEN, K.K. (2008): The invasive amphipod species *Gammarus tigrinus* (Sexton 1939) can rapidly change littoral communities in the Gulf of Finland (Baltic Sea). Aquatic Invasions 3: 405-412
- PERSSON, L.-E. (2001): Dispersal of *Platorchestia platensis* (Kröyer) (Amphipoda: Talitridae) along Swedish coasts: A slow but successful process. Estuar Coast Shelf Sci 52: 201-210
- PLATVOET, D., DE BRUYNE, R.H. & GMELIG MEYLING, A.W. (1995): Description of a new *Caprella*-Species from the Netherlands: *Caprella macho* nov. spec. (Crustacea, Amphipoda, Caprellidea). Bull Zool Mus 15: 1-4
- PLATVOET, D., DICK, J.T.A., MACNEIL, C., VAN RIEL, M.C. & VAN DER VELDE, G. (2009): Invader-invader interactions in relation to environmental heterogeneity leads to zonation of two invasive amphipods, *Dikerogammarus villosus* (Sowinsky) and *Gammarus tigrinus* Sexton: amphipod pilot species project (AMPIS) report 6. Biol Invasions 11: 2085-2093
- PÖCKL, M. (2009): Success of the invasive Ponto-Caspian amphipod *Dikerogammarus villosus* by life history traits and reproductive capacity. Biol Invasions 11: 2021-2041
- REICHERT, K. & BEERMANN, J. (2011): First record of the Atlantic gammaridean amphipod *Melita nitida* Smith, 1873 (Crustacea) from German waters (Kiel Canal). Aquatic Invasions 6: 103-108
- REWICZ T., GRABOWSKI M., MACNEIL C., BACELA-SPYCHALSKA, K. (2014): The profile of a 'perfect' invader – the case of killer shrimp, *Dikerogammarus villosus*. Aquatic Invasions 9: 267-288
- RUDOLPH, K. (1997): Zum Vorkommen des Flohkrebse *Pontogammarus robustoides* im Peenemündungsgebiet. Nat Mus 127: 306-312
- RÜPPELL, G. (1967): Zur Lokomotionsaktivität des Amphipoden *Orchestia platensis* im Freiland und im Laboratorium. Helgoländer Meeresunters 15: 172-180
- SCELLENBERG, A. (1942): Die Tierwelt Deutschlands und der angrenzenden Meeresteile, 40. Teil, IV: Flohkrebse oder Amphipoda, 252 pp
- SCHMITZ, W. (1960): Die Einbürgerung von *Gammarus tigrinus* Sexton auf dem europäischen Kontinent. Arch Hydrobiol 57: 223-225
- SCHREY, I. (2006): Verbreitung des eingeschleppten Gespensterkrebse *Caprella mutica* in der deutschen Bucht und seine potentielle Ansiedlung in natürlichen Habitaten. Diplomarbeit, Universität Rostock, 78 pp

- SCHÜCKEL, U., SCHÜCKEL, S., BECK, M. & LIEBEZEIT, G. (2010): New range expansion of *Caprella mutica* Schurin, 1935 (Malacostraca: Caprellidae) to the German coast, North Sea. *Aquatic Invasions* 5, Suppl 1: S85-S89
- SCHÜLER, L., SCHANZ, A., WINKELMANN, C., GRAF, S. & SCHMIDT, A. (2013): *Jassa* spp. (Amphipoda: Ischyroceridae) in der Ostsee – natürliche Arealerweiterung oder anthropogenes Phänomen? Oral presentation (and abstract), 16. Crustaceologentagung, E.-M.-A.-Universität Greifswald, März 2013
- SPICER, J.I. & JANAS, U. (2006): The beachflea *Platorchestia platensis* (Krøyer, 1845): a new addition to the Polish fauna (with a key to Baltic talitrid amphipods). *Oceanologia* 48: 287-295
- STRODE, E., BEREZINA, N.A., KALNINS, M. & BALODE, M. (2013): New records of the amphipods *Gammarus tigrinus* Sexton, 1939 and *Pontogammarus robustoides* G.O. Sars, 1894 in Latvian waters of the Baltic Sea. *BiolInvasions Records* 2: 63-68
- SZANIAWSKA, A., ŁAPUCKI, T. & NORMANT, M. (2003): The invasive amphipod *Gammarus tigrinus* Sexton, 1939, in Puck Bay. *Oceanologia* 45: 507-510
- TEIGSMARK, G. (1981): *Orchestia platensis* (Talitridae), an amphipod new to the Norwegian fauna. *Sarsia* 66: 165
- WAWRZYŃIAK-WYDROWSKA, B. & GRUSZKA, P. (2005): Population dynamics of alien gammarid species in the River Odra estuary. *Hydrobiologia*, 539:13-25
- WILDISH, D.J. & LINCOLN, R.J. (1979): Occurrence of *Orchestia platensis* Krøyer, 1845 (Amphipoda, Talitridae) in Britain. *Crustaceana* 36: 199-200
- WOUTERS, K. (2002): On the distribution of alien non-marine and estuarine macrocrustaceans in Belgium. *Bull koninklijk Belg Inst Natuurwetenschap Biol* 72: 119-129
- ZAIKO, A. & OLENIN, S. (2004): Impact of invasive benthic crustaceans on the resuspension of bottom sediments: an experimental study approach. *Logic and Hydrobiol Stud* 33 (3): 99-110
- ZETTLER, M.L. (1998): Zur Verbreitung der Malacostraca (Crustacea) in den Binnen- und Küstengewässern von Mecklenburg-Vorpommern. *Lauterbornia* 32: 49-65
- ZETTLER, M.L. (1999): Untersuchungen zum Makrozoobenthos des Breitlings (südliche Ostsee) unter besonderer Berücksichtigung der Crustacea. *Rostock Meeresbiol Beitr* 7: 79-90
- ZETTLER, M.L. (2008): Veränderung einer litoralen Amphipoda-Gemeinschaft am Beispiel einer Langzeitstudie im Oderhaff. *Lauterbornia* 62: 27-32
- ZETTLER M. (2015): Kurze Notiz über die Ankunft von *Echinogammarus trichiatus* im Ostseegebiet und den Erstnachweis von *Paramysis lacustris* in Deutschland. *Lauterbornia* 79: 151-156

- ANGER, K. (1990): Der Lebenszyklus der Chinesischen Wollhandkrabbe (*Eriocheir sinensis*) in Norddeutschland: Gegenwärtiger Stand des Wissens und neue Untersuchungen. Seevögel 11: 32-37
- ASAKURA, A. & WATANABE, S. (2005): *Hemigrapsus takanoi*, new species, a sibling species of the common Japanese intertidal crab *H. penicillatus* (Decapoda: Brachyura: Grapsoidea). J Crust Biol 25: 279-292
- ASHELBY, C.W., WORSFOLD, T.M. & FRANSEN, H.J.M. (2004): First records of the oriental prawn *Palaemon macrodactylus* (Decapoda: Caridea), an alien species in European waters, with a revised key to British Palaemonidae. J Mar Biol Ass U.K. 84: 1041-1050
- ASHELBY, C.W., DE GRAVE, S. & JOHNSON, M.L. (2013): The global invader *Palaemon macrodactylus* (Decapoda, Palaemonidae): An interrogation of records and a synthesis of data. Crustaceana 86: 594-624
- BACEVIČIUS, E. & GASIŪNAITĖ, Z.R. (2008): Two crab species – Chinese mitten crab (*Eriocheir sinensis* Edw.) and mud crab (*Rhitropanopeus harrisi* (Gould) ssp. *tridentatus* (Maitland)) in the Lithuanian coastal waters, Baltic Sea. Transit Waters Bull 2: 63-68
- BÉGUER, M., GIRARDIN, M. & BOËT, P. (2007): First record of the invasive oriental shrimp *Palaemon macrodactylus* Rathbun, 1902 in France (Gironde Estuary). Aquatic Invasions 2: 132-136
- BÉGUER, M., BERGÉ, J., MARTIN, J., MARTINET, J., PAULIAC, G., GIRARDIN, M. & BOËT, P. (2011): Presence of *Palaemon macrodactylus* in a European estuary: evidence for a successful invasion of the Gironde (SW France). Aquatic Invasions 6: 301-318
- BOETS, P., LOCK, K. & GOETHALS, L.M. (2012): Assessing the importance of alien macrocrustacea (Malacostraca) within macroinvertebrate assemblages in Belgian coastal harbours. Helgol Mar Res 66: 175-187
- BOETTGER, C.R. (1933b): Die Ausbreitung der Wollhandkrabbe (*Eriocheir sinensis* Milne-Edw.) in Europa. Sitzungsber Ges naturforsch Freunde v Berlin: 399-415
- BRETON, G., FAASSE, M., NOË, I.P. & VINCENT, T. (2002): A new alien crab in Europe: *Hemigrapsus sanguineus* (Decapoda: Brachyura: Grapsidae). Journal of Crustacean Biology 22: 184-189
- CHÍCARO, M.A., LEITÃO, T., RANGE, P., GUTIERREZ, C., MORALES, J., MORAIS, P. & CHÍCARO L. (2009): Alien species in the Guadiana Estuary (SE-Portugal/SW-Spain): *Blackfordia virginica* (Cnidaria, Hydrozoa) and *Palaemon macrodactylus* (Crustacea, Decapoda): potential impacts and mitigation measures. Aquatic Invasions 4: 501-506
- CUESTA, J.A., GONZÁLEZ-ORTEGÓN, E., DRAKE, P. & RODRÍGUEZ, A. (2004): First record of *Palaemon macrodactylus* Rathbun, 1902 (Decapoda, Caridea, Palaemonidae) from European waters. Crustaceana 77: 377-380
- DAUVIN, J.-C. (2009a): Establishment of the invasive Asian shore crab *Hemigrapsus sanguineus* (De Haan, 1835) (Crustacea: Brachyura: Grapsoidea) from the Cotentin Peninsular, Normandy, France. Aquatic Invasions 4: 467-472

- DAUVIN, J.-C. (2009b): Asian Shore Crabs *Hemigrapsus* spp. (Crustacea: Brachyura: Grapsoidea) continue their invasion around the Cotentin Peninsula, Normandy, France: Status of the *Hemigrapsus* population in 2009. *Aquatic Invasions* 4: 605-611
- DAUVIN, J.-C., RIUS, A.T. & RUELLET, T. (2009): Recent expansion of two invasive crabs species *Hemigrapsus sanguineus* (de Haan, 1835) and *H. takanoi* Asakura and Watanabe 2005 along the Opal Coast, France. *Aquatic Invasions* 4: 451-465
- DAUVIN, J.-C. & DELHAY, J.-B. (2010): First record of *Hemigrapsus takanoi* (Crustacea: Decapoda: Grapsidae) on the western coast of northern Cotentin, Normandy, western English Channel. *Mar Biodivers Rec* 3: 1-3
- DAUVIN, J.-C. & DUFOSSÉ, F. (2011): *Hemigrapsus sanguineus* (De Haan, 1835) (Crustacea: Brachyura: Grapsoidea) a new invasive species in European waters: the case of the French English Channel coast (2008-2010). *Aquatic Invasions* 6: 429-438
- DÜRR, A. (2010): Untersuchungen zur Populations- und Fortpflanzungsbiologie von Garnelen in der westlichen Ostsee. Diplomarbeit, Universität Kiel, 160 pp
- FLAUNDERS, E. (2000): Untersuchungen zur Bestandsregulierung und Verwertung der Chinesischen Wollhandkrabbe (*Eriocheir sinensis*) unter besonderer Berücksichtigung der Fischereiverhältnisse im Elbe/Havel-Gebiet. Schriften des Instituts für Binnenfischerei e. V. Potsdam-Sacrow, Bd. 5, 82 pp
- FOWLER, A.E., FORSSTRÖM, T., VON NUMERS, M. & VESAKOSKI, O. (2013): The North American mud crab *Rhithropanopeus harrisii* (Gould, 1841) in newly colonized Northern Baltic Sea: distribution and ecology. *Aquatic Invasions* 8: 89-96
- GAGNON, K. & BOSTRÖM, C. (2016): Habitat expansion of the Harris mud crab *Rhithropanopeus harrisii* (Gould, 1841) in the northern Baltic Sea: potential consequences for the eelgrass food web. *BiolInvasions Rec* 5: 101-106
- GEBURZI, J.C., GRAUMANN, G., KÖHNK, S. & BRANDIS, D. (2015): First record of the Asian crab *Hemigrapsus takanoi* Asakura & Watanabe, 2005 (Decapoda, Brachyura, Varunidae) in the Baltic Sea. *BiolInvasions Rec* 4: 103-107
- GERARD, V.A., CERRATO, R.M. & LARSON, A.A. (1999): Potential impacts of a western Pacific grapsid crab on intertidal communities of the northwestern Atlantic Ocean. *Biol Invasions* 1: 353-361
- GILBEY, V., ATTRILL, M.J. & COLEMAN, R.A. (2008): Juvenile Chinese mitten crabs (*Eriocheir sinensis*) in the Thames estuary: distribution, movement and possible interactions with the native crab *Carcinus maenas*. *Biol Invasions* 10: 67-77
- GOLLASCH, S. (1999): The Asian decapod *Hemigrapsus penicillatus* (de Haan, 1835) (Grapsoidea, Decapoda) introduced in European waters: *status quo* and future perspective. *Helgoländer Meeresunters* 52: 359-366
- GOLLASCH, S. (2011): NOBANIS – Invasive Alien Species Fact Sheet – *Eriocheir sinensis*. From: Online database of the North European and Baltic Network on Invasive Alien Species – NOBANIS www.nobanis.org (10.07.2013)
- GONZÁLEZ-ORTEGÓN, E. & CUESTA, J.A. (2006): An illustrated key to species of *Palaemon* and *Palaemonetes* (Crustacea: Decapoda: Caridea) from European waters, including the alien species *Palaemon macrodactylus*. *J Mar Biol Ass U.K.* 86: 93-102

- GONZÁLEZ-ORTEGÓN, E., CUESTA, J.A. & SCHUBART, C.D. (2007): First report of the oriental shrimp *Palaemon macrodactylus* Rathbun, 1902 (Decapoda, Caridea, Palaemonidae) from German waters. *Helgol Mar Res* 61: 67–69
- GONZÁLES-ORTEGÓN, E., CUESTA, J.A., PASCUAL, E. & DRAKE, P. (2010): Assessment of the interaction between the white shrimp, *Palaemon longirostris*, and the exotic oriental shrimp, *Palaemon macrodactylus*, in a European estuary (SW Spain). *Biol Invasions* 12: 1731-1745
- GOTHLAND, M., DAUVIN, J.-C., DENIS, L., JOBERT, S., OVAERT, J., PEZY, J.-P. & SPILMONT, N. (2013): Additional records and distribution (2011-2012) of *Hemigrapsus sanguineus* (De Haan, 1835) along the French coast of the English Channel. *Management Biol Invasions* 4: 305-315
- HÄNFLING, B., CARVALHO, G.R. & BRANDL, R. (2002): mt-DNA sequences and possible invasion pathways of the Chinese mitten crab. *Mar Ecol Prog Ser* 238: 307-310
- HERBORG, L.-M., RUSHTON, S.P., CLARE, A.S. & BENTLEY, M.G. (2003): Spread of the Chinese mitten crab (*Eriocheir sinensis* H. Milne Edwards) in continental Europe: analysis of a historical data set. *Hydrobiologia* 503: 21-28
- HERBORG, L.-M., RUSHTON, S.P., CLARE, A.S. & BENTLEY, M.G. (2005): The invasion of the Chinese mitten crab (*Eriocheir sinensis*) in the United Kingdom and its comparison to continental Europe. *Biol Invasions* 7: 959-968
- JANAS, U. & TUTAK, B. (2014): First record of the oriental shrimp *Palaemon macrodactylus* M.J. Rathbun, 1902 in the Baltic Sea. *Oceanol Hydrobiol Stud* 43: 431-435
- JENSEN, K.R. (2010i): NOBANIS – Invasive Alien Species Fact Sheet – *Callinectes sapidus*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (09.07.2013)
- JENSEN, K.R. (2010j): NOBANIS – Invasive Alien Species Fact Sheet – *Rhithropanopeus harrisi*. From: Identification key to marine invasive species in Nordic waters. NOBANIS www.nobanis.org (30.07.2013)
- KINNE, O. & ROTTHAUWE, H.-W. (1952): Biologische Beobachtungen und Untersuchungen über die Blutkonzentration an *Heteropanope tridentatus* MAITLAND (Dekapoda). *Kieler Meeresforsch* 8: 212-217
- KOTTA, J. & OJAVEER, H. (2012): Rapid establishment of the alien crab *Rhithropanopeus harrisi* (Gould) in the Gulf of Riga. *Estonian J Ecol* 61: 293-298
- KÜHL, H. (1977a): *Mercierella enigmatica* (Polychaeta: Serpulidae) an der deutschen Nordseeküste. *Veröff Inst Meeresforsch Bremerh* 16: 99-104
- KÜHL, H. (1977b): Neue Tierarten an der deutschen Nordseeküste. *Inform Fischwirtsch* 24: 61-62
- LANDSCHOFF, J., LACKSCHEWITZ, D., KESY, K. & REISE, K. (2013): Globalization pressure and habitat change: Pacific rocky shore crabs invade armored shorelines in the Atlantic Wadden Sea. *Aquatic Invasions* 8: 77-87
- LAVESQUE, N., BACHELET, G., BÉGUER, M., GIRARDIN, M., LEPAGE, M., BLANCHET, H., SORBE, J.-C., MODÉLAN, J., SAURIAU, P.-G. & AUBY, I. (2010): Recent expansion of the oriental

- shrimp *Palaemon macrodactylus* (Crustacea: Decapoda) on the western coasts of France. *Aquatic Invasions* 5, Suppl: S103-S108
- LOHRER, A.M. & WHITLATCH, R.B. (2002): Interactions among aliens: apparent replacement of one exotic species by another. *Ecology* 83: 719-732
- MARKERT, A., RAUPACH, M., SEGELKEN-VOIGT, A. & WEHRMANN, A. (2014): Molecular identification and morphological characteristics of native and invasive Asian brush-clawed crabs (Crustacea: Brachyura) from Japanese and German coasts: *Hemigrapsus penicillatus* (De Haan, 1835) versus *Hemigrapsus takanoi* Asakura & Watanabe 2005. *Org Divers Evol* 14: 369-382
- MACDONALD, J.A., ROUDEZ, R., GLOVER, T. & WEIS, J.S. (2007): The invasive green crab and Japanese shore crab: behavioural interactions with a native crab species, the blue crab. *Biol Invasions* 9: 837-848
- MICU, D. & NIȚĂ, V. (2009): First record of the Asian prawn *Palaemon macrodactylus* Rathbun, 1902 (Caridea: Palaemonoidea: Palaemonidae) from the Black Sea. *Aquatic Invasions* 4: 597-604
- MICU, D., NIȚĂ, V. & TODOROVA, V. (2010): First record of the Japanese shore crab *Hemigrapsus sanguineus* (de Haan, 1835) (Brachyura: Grapsoidea: Varunidae) from the Black Sea. *Aquatic Invasions* 5, Suppl: S1-S4
- NEHRING, S. (2000b): Zur Bestandssituation von *Rhithropanopeus harrisi* (Gould, 1841) in deutschen Gewässern: Die sukzessive Ausbreitung eines amerikanischen Neozoons (Crustacea: Decapoda: Panopeidae). *Senckenbergiana marit* 30: 115-122
- NEHRING, S. & LEUCHS, H. (1999b): *Rhithropanopeus harrisi* (Gould, 1841) (Crustacea: Decapoda) – ein amerikanisches Neozoon im Elbeästuar. *Lauterbornia* 35: 49-51
- NEHRING, S., SPECKELS, G. & ALBERSMEYER, J. (2008): The American blue crab *Callinectes sapidus* RATHBUN on the German North Sea coast: Status quo and further perspectives. *Senckenbergiana marit* 38: 39-44
- NEHRING, S. & VAN DER MEER, U. (2010): First record of a fertilized female blue crab, *Callinectes sapidus* Rathbun, 1896 (Crustacea: Decapoda: Brachyura), from the German Wadden Sea and subsequent secondary prevention measures. *Aquatic Invasions* 5: 215-218
- NOËL, P., TARDY, E. & D'UDEKEM D'ACQZ, C. (1997): Will the crab *Hemigrapsus penicillatus* invade the coasts of Europe? *Comptes Rendus de l'Académie des Sciences, Série 3, Sciences de la vie*, 320: 741-745
- OBERT, B., HERLYN, M. & GROTHJAHN, M. (2007): First records of two crabs from the North West Pacific *Hemigrapsus sanguineus* und *H. takanoi* at the coast of Lower Saxony, Germany. *Wadden Sea Newsletter* 1: 21-22
- O'CONNOR, N.J. (2014): Invasion dynamics on a temperate rocky shore: from early invasion to establishment of a marine invader. *Biol Invasions* 16: 73-87
- OJAVEER, H., GOLLASCH, S., JAANUS, A., KOTTA, J., LAINE, A.O., MINDE, A., NORMANT, M. & PANOV, V.E. (2007): Chinese mitten crab *Eriocheir sinensis* in the Baltic Sea – a supply-side invader? *Biol Invasions* 9: 409-418

- OTTO, T. & BRANDIS, D. (2011): First evidence of *Eriocheir sinensis* reproduction from Schleswig-Holstein, Northern Germany, western Baltic Sea. *Aquatic Invasions* 6, Suppl: S65-S69
- PETERS, N. (1938): Ausbreitung und Verbreitung der Chinesischen Wollhandkrabbe (*Eriocheir sinensis* M. Edw.) in Europa in den Jahren 1933 bis 1935. *Mitt Hamb Zool Mus Inst* 47: 1-31
- PETERS, N., PANNING, A. & SCHNAKENBECK (1933): Die Chinesische Wollhandkrabbe (*Eriocheir sinensis*) in Deutschland. *Zool Anz, Ergänzungsband zu Bd.* 104
- PEZY, J.-P., DAUVIN, J.-C. & VINCENT, T. (2015): New records of *Eriocheir sinensis* H. Milne Edwards, 1853 in Normandy. *Mar Biodiv Rec* 8, e15, 1-4, publ online
- RAYKOV, V.ST., LEPAGE, M. & PÉREZ-DOMÍNGUEZ (2010): First record of oriental shrimp, *Palaemon macrodactylus* Rathbun, 1902 in Varna Lake, Bulgaria. *Aquatic Invasions* 5, Suppl 1: S91-S95
- ROCHE, D.G. & TORCHIN, M.E. (2007): Established population of the North American Harris mud crab, *Rhithropanopeus harrisi* (Gould 1841) (Crustacea: Brachyura: Xanthidae), in the Panama Canal. *Aquatic Invasions* 2: 155-161
- ROUDEZ, R.J., GLOVER, T. & WEIS, J.S. (2008): Learning in an invasive and a native predatory crab. *Biol Invasions* 10: 1191-1196
- RUIJTER, R. de (2008): cs-verslag. *Zeepard* 68: 2-7
- SCHUBART, C.D. (2003): The East Asian shore crab *Hemigrapsus sanguineus* (Brachyura: Varunidae) in the Mediterranean Sea: An independent human-mediated introduction. *Sci Mar* 67: 195-200
- SCHÜCKEL, U., MARKERT, A., NEUMANN, H., KRÖNCKE, I. & WEHRMANN, A. (2013): Neue Krebse im Vormarsch. *Senckenberg Natur, Forschung, Museum* 143: 152-157
- SCHULZ, R. & SMETANA, P. (2001): Occurrence of native and introduced crayfish in Northeastern Germany and Northwestern Poland. *Bull Fr Pêche Piscic* 361: 629-641
- SEELEY, B., SEWELL, J. & CLARK, P.F. (2015): First GB records of the invasive Asian shore crab, *Hemigrapsus sanguineus* from Glamorgan, Wales and Kent, England. *Mar Biodivers Rec* 8; e102; published online
- SOORS, J., FAASSE, M., STEVENS, M., VERBESSEM, I., DE REGGE, N. & VAN DEN BERGH, E. (2010): New crustacean invaders in the Schelde estuary (Belgium). *Belg J Zool* 140: 3-10
- SPIVAK, E.D., BOSCHI, E.E. & MARTORELLI, S.R. (2006). Presence of *Palaemon macrodactylus* Rathbun, 1902 (Crustacea: Decapoda: Caridea: Palaemonidae) in Mar del Plata harbor, Argentina: first record from southwestern Atlantic waters. *Biol Invasions* 8: 673-676
- STEBBING, P., JOHNSON, P., DELAHUNTY, A., CLARK, P.F., MCCOLLIN, T., HALE, C. & CLARK, S. (2012): Reports of American lobsters, *Homarus americanus* (H. Milne Edwards, 1837), in British waters. *BiolInvasions Records* 1: 17-23

- D'UDEKEM D'ACUZ, C., FAASSE, M., DUMOULIN, E. & DE BLAUWE, H. (2005): Occurrence of the Asian shrimp *Palaemon macrodactylus* in the southern bight of the North Sea, with a key to the Palaemonidae of north-western Europe (Crustacea: Decapoda: Caridea). *Ned Faun Meded* 22: 95-112
- VAN DER MEEREN, G., STØTTRUP, J., ULMESTRAND, M., ØRESLAND, V., KNUTSEN, J.A. & AGNALT, A.-L. (2010): NOBANIS – Invasive Alien Species Fact Sheet – *Homarus americanus*. – From: Online Database of the European Network on Invasive Alien Species. NOBANIS www.nobanis.org (18.07.2014)
- WESTMAN, K. (2002): Alien crayfish in Europe: Negative and positive impacts and interactions with native crayfish. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): *Invasive aquatic species of Europe. Distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, Boston, London, 76-95
- WITTFOTH, A.K.J. & ZETTLER, M.L. (2013): The application of a Biopollution Index in German Baltic estuarine and lagoon waters. *Management of Biol Invasions* 4: 43-50
- WOOD, C.A., BISHOP, J.D.D., DAVIES, C.J., DELDUCA, E.L., HATTON, J.C., HERBERT, R.J.H. & CLARK, P.F. (2015): *Hemigrapsus takanoi* Asakura and Watanabe, 2005 (Crustacea: Decapoda: Brachyura: Grapsoidea): First records of the brush-clawed shore crab from Great Britain. *BiolInvasions Rec* 4: 109-113
- WORSFOLD, T.M. & ASHELBY, C.W. (2006): Additional UK records of the non-native prawn *Palaemon macrodactylus* (Crustacea: Decapoda). *Mar Biodivers Rec* 1: 1-3
- WOUTERS, K. (2002): On the distribution of alien non-marine and estuarine macrocrustaceans in Belgium. *Bull koninklijk Belg Inst Natuurwetenschap Biol* 72: 119-129
- ZETTLER, M.L. (1998): Zur Verbreitung der Malacostraca (Crustacea) in den Binnen- und Küstengewässern von Mecklenburg-Vorpommern. *Lauterbornia* 32: 49-65
- ZETTLER, M. (2001): Die höheren Krebse (Malacostraca) der Darß-Zingster Boddenkette. *Meer Mus* 16: 74-75

ÜBERSICHTSARTIKEL UND LISTEN MIT ANGABEN ZU CRUSTACEA

- ENO, N.C., CLARK, R.A. & SANDERSON, W.G. (eds.) (1997): *Non-native marine species in British waters: a review and directory*. Joint Nature Conservation Committee, Peterborough, 152 pp
- FRANKE, H.-D. & GUTOW, L. (2004): Long-term changes in the macrobenthos around the rocky island of Helgoland (German Bight, North Sea). *Helgol Mar Res* 58: 303-310
- GERLACH, S.A. (2000): *Die Biodiversität in der deutschen Nord- und Ostsee, Bd. 1, Checkliste der Fauna der Kieler Bucht und eine Bibliographie zur Biologie und Ökologie der Kieler Bucht*. Bundesanstalt für Gewässerkunde, Koblenz, 376 pp
- GITTENBERGER, A., RENSING, M., STEGENGA, H. & HOEKSEMA, B.W. (2010): Native and non-native species of hard substrata in the Dutch Wadden Sea. *Ned Faun Meded* 33: 21-76

- GOLLASCH, S. & MECKE, R. (1996): Eingeschleppte Organismen. In: J.L. LOZAN, R. LAMPE, W. MATTHAUS, E. RACHOR, H. RUMOHR & H. V. WESTERNHAGEN (Eds.): Warnsignale aus der Ostsee. Parey Buchverlag, Berlin: 146-150
- GOLLASCH, S. & NEHRING, S. (2006): National checklist for aquatic species in Germany
Aquatic invasions 1: 245–269
- GOLLASCH, S., HAYDAR, D., MINCHIN, D., WOLFF, W.J. & REISE, K. (2009): Introduced Aquatic Species of the North Sea Coasts and Adjacent Brackish Waters. In: RILOV, G. & CROOKS, J.A. (eds.): Biological invasions in marine ecosystems, Chapter 29: 507-528
Ecological Studies 204, Springer-Verlag Berlin Heidelberg 2008
- GOMOIU, M.-T., ALEXANDROV, B., SHADRIN, N. & ZAITSEV, Y. (2002): The Black Sea – a recipient, donor and transit area for alien species. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): Invasive aquatic species of Europe. Distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, Boston, London, 341-350
- HAYWARD, P.J. & RYLAND, J.S. (1990): The Marine fauna of the British Isles and North-West Europe, Vol. 1 & 2. Oxford Science Publications, Clarendon Press, Oxford, 1990
- HOPKINS, C.C.E. (2002): Introduced marine organisms in Norwegian waters, including Svalbard, In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): Invasive aquatic species of Europe. Distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, Boston, London, 240-252
- JENSEN, K. & KNUDSEN, J. (2005): A summary of alien marine benthic invertebrates in Danish waters. *Oceanol Hydrobiol Stud* 34, Suppl 1: 137-162
- KERCKHOF, F., HAELTERS, J. & GOLLASCH, S. (2007): Alien species in the marine and brackish water ecosystems: the situation in Belgian waters. *Aquatic Invasions* 2: 243-257
- LACKSCHEWITZ, D., REISE, K. & BUSCHBAUM, C. (2009, 2010, 2011): Schnellerfassung von Neobiota in deutschen Küstengewässern. Im Auftrag des Landesamtes für Landwirtschaft, Umwelt und ländliche Räume des Landes Schleswig-Holstein
- LACKSCHEWITZ, D., REISE, K. & BUSCHBAUM, C. (2012, 2013): Neobiota. In: Monitoring und Bewertung des Benthos, der Lebensraumtypen/Biotope und der Gebietsfremden Arten (Cluster 4, Benthosmonitoring, Synthetischer Jahresbericht)
- LEPPÄKOSKI, E. (1984): Introduced species in the Baltic Sea. *Ophelia*, Suppl 3: 123-135
- LEPPÄKOSKI, E. & OLENIN, S. (2000): Non-native species and rates of spread: lessons from the brackish Baltic Sea. *Biol Invasions* 2: 151-163
- LEPPÄKOSKI, E., OLENIN, S. & GOLLASCH, S. (2002): The Baltic Sea – a field laboratory for invasion biology. In: LEPPAKOSKI, E., GOLLASCH, S. & OLENIN, S. (eds.): Invasive aquatic species of Europe. Distribution, impacts and management. Kluwer Academic Publishers, Dordrecht, Boston, London, 253-259
- LINCOLN, R.J. (1979): British marine amphipoda: Gammaridea. British Museum (Natural History), 658 pp
- MINCHIN, D., COOK, E.J. & CLARK, P.F. (2013): Alien species in British brackish and marine waters. *Aquatic Invasions* 8, Suppl: 3-19

- MÜLLER, R. & PESCHEL, T. (2007): Eingebürgerte Arten des Makrozoobenthos und der submersen natanten Makrophyten in Berliner Gewässern. Planungsbüro Hydrobiologie Berlin, im Auftrag der Senatsverwaltung für Stadtentwicklung Berlin, 44 pp
- NEHRING, S. & LEUCHS, H. (1999a): Neozoa (Makrobenthos) an der deutschen Nordseeküste. Eine Übersicht. Bundesanstalt für Gewässerkunde, Koblenz. 131 pp
- OSTERKAMP, S. & SCHIRMER, M. (2000): Klimasensitivität der Unterweser und ihrer Vorländer. In: Bund-Länder-Programm ‚Klimaänderung und Küste‘, Projekt ‚Klimaänderung und Unterweserregion‘ (KLIMU) (Fallstudie Weserästuar); Abschlussbericht Teilprojekt Ökologischer Komplex, Universität Bremen, 151 pp
- PANOV, V.E., KRYLOV, P.I. & TELESH, I.V. (1999): The St. Petersburg harbour profile. In: GOLLASCH, S. & LEPPÄKOSKI, E. (eds.) Initial risk assessment of alien species in Nordic coastal waters. Nord 1999:8. Nordic Council of Ministers, Copenhagen, Denmark: 225-244
- POST, D. & LANDMANN, M. (1994): Verbreitungsatlas der Fließgewässerfauna in Ostfriesland. Staatl Amt Wasser Abfall, 141 S.
- SCHÖLL, F., BECKER, C. & TITTIZER, T. (1995): Das Makrozoobenthos des schiffbaren Rheins von Basel bis Emmerich 1986-1995. *Lauterbornia* 21: 115-137
- TITTIZER, T. (1997): Ausbreitung aquatischer Neozoen (Makrobenthos) in den europäischen Wasserstraßen, erläutert am Beispiel des Main-Donau-Kanals Schriftenreihe des Bundesamtes f. Wasserwirtschaft 4: 113-134
- TITTIZER, T., SCHÖLL, F., BANNING, M., HAYBACH, A. & SCHLEUTER, M. (2000): Aquatische Neozoen im Makrobenthos der Binnenwasserstraßen Deutschlands. *Lauterbornia* 39: 1-72
- WASMUND, N., POLLEHNE, F., POSTEL, L., SIEGEL, H. & ZETTLER, M.L. (2007): Biologische Zustandseinschätzung der Ostsee im Jahre 2006. *Meereswiss Ber, Warnemünde*, 71: 1-82
- WASMUND, N., POSTEL, L. & ZETTLER, M.L. (2012): Biologische Bedingungen in der deutschen ausschließlichen Wirtschaftszone der Nordsee im Jahre 2011. *Meereswiss Ber, Warnemünde*, 90
- WEIDEMA, I.R. (2000): Introduced species in the Nordic Countries, *Nord*, 13: 242 pp
- WITT, J. & KRUMWIEDE, A. (2006): Entwicklung eines Bewertungskonzeptes und Bewertung des Makrozoobenthos der Unterweser. Anhang: Beschreibung und Bewertung des Ist-Zustandes. Im Auftrag des Wasser- und Schifffahrtsamtes Bremerhaven (WSA)
- WOLFF, W.J. (2005): Non-indigenous marine and estuarine species in The Netherlands. *Zool Meded* 79: 1-116
- Zettler, M.L. & Röhner, M. (2004): Die Biodiversität in der deutschen Nord- und Ostsee, Bd. 3: Verbreitung und Entwicklung des Makrozoobenthos der Ostsee zwischen Fehmarnbelt und Usedom – Daten von 1839 bis 2001. BfG-1421

Listen und Internetquellen zu Crustacea

ICES WGITMO Report 2005

www.arge-elbe.de/wge/Download/Texte/02Amph.pdf

www.ciesm.org (The Mediterranean Science Commission)

www.corpi.ku.lt/nemo/alien_species_directory.html (Baltic Sea Alien Species Database)

www.europe-aliens.org/index.jsp (DAISIE Datenbank)

www.frammandearter.se

www.natuurbericht.nl

www.nobanis.org

www.sea.ee

www.vattenfall.dk/da/file/review-report-2005